

ASFMR1 splice variant

A predictor of fragile X-associated tremor/ataxia syndrome

Padmaja Vittal, MD, MS, Shrikant Pandya, BS, Kevin Sharp, MS, Elizabeth Berry-Kravis, MD, PhD, Lili Zhou, MD, Bichun Ouyang, PhD, Jonathan Jackson, BS, and Deborah A. Hall, MD, PhD

Correspondence

Dr. Hall
deborah_a_hall@rush.edu

Neurol Genet 2018;4:e246. doi:10.1212/NXG.0000000000000246

Abstract

Objective

To explore the association of a splice variant of the *antisense fragile X mental retardation 1* (*ASFMR1*) gene, loss of *fragile X mental retardation 1* (*FMR1*) AGG interspersions and *FMR1* CGG repeat size with manifestation, and severity of clinical symptoms of fragile X-associated tremor/ataxia syndrome (FXTAS).

Methods

Premutation carriers (PMCs) with FXTAS, without FXTAS, and normal controls (NCs) had a neurologic evaluation and collection of skin and blood samples. Expression of *ASFMR1* transcript/splice variant 2 (*ASFMR1-TV2*), nonspliced *ASFMR1*, total *ASFMR1*, and *FMR1* messenger RNA were quantified and compared using analysis of variance. Least absolute shrinkage and selection operator (LASSO) logistic regression and receiver operating characteristic analyses were performed.

Results

Premutation men and women both with and without FXTAS had higher *ASFMR1-TV2* levels compared with NC men and women ($n = 135,135$, $p < 0.0001$), and *ASFMR1-TV2* had good discriminating power for FXTAS compared with NCs but not for FXTAS from PMC. After adjusting for age, loss of AGG, larger CGG repeat size (in men), and elevated *ASFMR1-TV2* level (in women) were strongly associated with FXTAS compared with NC and PMC (combined).

Conclusions

This study found elevated levels of *ASFMR1-TV2* and loss of AGG interruptions in both men and women with FXTAS. Future studies will be needed to determine whether these variables can provide useful diagnostic or predictive information.

From the Department of Neurosciences (P.V.), Northwestern Medicine Regional Medical Group, Winfield, IL; the Department of Bioengineering (S.P.), University of Illinois at Chicago; the Department of Pediatrics, Neurological Sciences, Biochemistry (K.S., E.B.-K., L.Z., J.J.), and the Department of Neurological Sciences (B.O., D.A.H.), Rush University Medical Center, Chicago, IL.

Funding information and disclosures are provided at the end of the article. Full disclosure form information provided by the authors is available with the full text of this article at Neurology.org/NG.

The Article Processing Charge was paid for by *Neurology: Genetics*.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

AR = activation ratio; **ASFMR1** = antisense fragile-X mental retardation 1; **ANOVA** = analysis of variance; **cDNA** = complementary DNA; **FMRI** = fragile X mental retardation 1; **FXS** = fragile X syndrome; **FXTAS** = fragile X-associated tremor/ataxia syndrome; **FXTAS-RS** = FXTAS Rating Scale; **LASSO** = least absolute shrinkage and selection operator; **mRNA** = messenger RNA; **NC** = normal control; **PMC** = premutation carrier; **ROC** = receiver operating characteristic; **TV2** = transcript variant 2; **UTR** = untranslated.

Fragile X-associated tremor/ataxia syndrome (FXTAS) is an inherited neurodegenerative disorder in adults older than 50 years because of 55–200 (permutation range) CGG repeat expansion in the 5' untranslated (UTR) region *fragile X mental retardation 1* (*FMRI*) gene causing adult-onset ataxia.¹ Primary features include progressive action tremor and cerebellar ataxia, whereas associated findings include parkinsonism, neuropathy, cognitive decline, and dysautonomia.² The CGG repeat in the 5' UTR is unstable and its repeat length in the normal population ranges from 6 to 55 repeats.^{3,4} More than 200 repeats (full mutation) is associated with methylation and transcriptional silencing of *FMRI* and consequent absence or deficiency of fragile X mental retardation protein, resulting in fragile X syndrome⁵ (FXS). The estimated prevalence of the *FMRI* premutation is approximately 1/430 males and 1/209 females.⁶ The molecular mechanism of FXTAS includes elevated levels of the expanded CGG repeat-containing messenger RNA (mRNA) transcript and abnormal repeat-associated non-AUG-initiated (RAN) translation products that lead to cellular toxicity and neuronal intranuclear inclusions (including DNA damage response proteins).^{7–9}

Although there is a relationship in a large group analysis between the age at onset/motor severity of FXTAS and CGG repeat size¹⁰ with increasing risk and severity at higher repeat lengths, the CGG repeat size alone does not fully explain risk. Women have a protective normal X allele, and hence, the propensity for developing FXTAS is different between men and women. In carrier women, a higher activation ratio (AR) (percentage of cells expressing the normal *FMRI* gene) may lower the presence and severity of motor signs of FXTAS.^{10–12} It is probable, however, that there are additional secondary genes, medical conditions, and environmental factors that affect the effects of the premutation expansion and contribute to the risk of developing symptoms.

A second gene, *antisense FMRI* (*ASFMR1*) overlaps a portion of the *FMRI* gene including the CGG repeat sequence, is transcribed in the reverse direction from *FMRI* with 5 splice variants of unclear exact clinical impact.¹³ One of the variants previously described¹³ *ASFMR1* splice variant 2 (*ASFMR1* transcript variant 2 [*ASFMR1-TV2*]) was further explored in this study. Elevated levels of the expanded GCC repeat-containing *ASFMR1* transcript are observed in premutation carriers (PMCs),¹³ and small unpublished studies have suggested that certain splice variants of the *ASFMR1* transcript

are associated with the presence of neurologic symptoms in PMC men with FXTAS. Within the CGG repeat element of the *FMRI* gene, there are AGG trinucleotide interruptions (typically separated by 9–11 CGG repeats), which are known to disrupt the otherwise pure CGG repeat motif. Normal *FMRI* alleles typically possess 2 or 3 AGG interruptions; premutation alleles generally possess 2 or less interruptions, whereas larger premutation alleles tend to have fewer AGG interruptions. The loss of AGG interruptions is believed to increase the probability that large normal and small premutation size alleles will be unstable on transmission and expand to a full mutation allele on transmission from small or moderate size alleles.¹⁴ The rationale for investigating AGG interspersions is that pilot data from our group show that a lack of AGG interspersions is associated with neurologic signs. In this study, we hypothesized that elevated specific splice variants (*ASFMR1-TV2*) of the *ASFMR1* transcript and a lack of AGG interspersions may play a role in FXTAS, and we attempted to determine the correlation between these features in development of clinical symptoms in FXTAS.

Methods

Participants

This is a case-control study. Adult PMCs and healthy NCs were recruited from the Rush University Fragile X-Associated Disorders Program. Recruitment occurred from 3 studies investigating neurologic phenotypes in PMCs and NCs^{15–17} through clinics, and fragile X community events. The published criteria for FXTAS were followed to classify as possible, probable, or definite FXTAS participants.^{3,4}

Standard protocol approvals, registrations, and patient consents

This study received approval from the Institutional Review Board at the Rush University Medical Center. Written informed consent was obtained from all patients (or guardians of patients) participating in the study. The family history of FXS from participants (but not FXTAS or other neurologic conditions in the family) was obtained.

Neurologic testing

Each subject was evaluated with a neurologic examination, the FXTAS Rating Scale (FXTAS-RS),¹⁸ and the Total Neuropathy Scale¹⁹ (modified to exclude those items that require a nerve conduction study). The FXTAS-RS was developed to measure the salient features of the movement disorder seen in

FXTAS. The scale was constructed by combining 3 published rating scales commonly used to assess tremor (Clinical Rating Scale for Tremor),²⁰ ataxia (International Cooperative Ataxia Rating Scale),²¹ parkinsonism (Unified Parkinson's Disease Rating Scale),²² and a tandem gait item (Huntington's Disease Rating Scale).²³

CGG and AGG determination

The *FMR1* PCR assay (Asuragen Inc, Austin, TX) was performed on blood samples from participants using a recently developed highly sensitive method, which detects all *FMR1* expansions and allows accurate quantification of allele-specific CGG repeat length,^{24,25} including identification of AGG interspersions, using primers that flank the repeat and internal repeat primers. Southern blot assay was performed on samples from premutation women to determine the normal AR.^{26,27}

Fibroblast collection

Skin biopsy was performed under sterile conditions and delivered to the laboratory immediately for processing. The skin biopsy was placed in culture media and later diced under sterile conditions on a culture plate and then plated in T25 flasks in CHANG-Amnio™-Medium (Irvine Scientific, Santa Ana, CA). The flasks were incubated at 37°C with 5% carbon dioxide. After cells grew out, longer-term cultures were maintained in Corning™ cellgro™ Cell Culture Media–Dulbecco's Modified Eagle's Medium supplemented with 10% fetal bovine serum (Gibco) and 1% Gibco® Antibiotic-Antimycotic (containing 10,000 units/mL of penicillin, 10,000 µg/mL of streptomycin, and 25 µg/mL of Gibco Amphotericin B). Aliquots of all cell lines were frozen and kept at –80° for long-term use in future studies.

ASFMR1 and *FMR1* messenger RNA assays

RNA was purified using the Qiagen RNeasy kit from blood or fibroblast samples. Samples were treated with DNase and screened for contamination; then, complementary DNA (cDNA) synthesis was performed using the High Capacity cDNA Reverse Transcriptase Kit from Applied Biosystems, Foster City, CA. The defining characteristic of *ASFMR1-TV2* is the 84-nucleotide deletion near the beginning of the open reading frame and does include the repeat. The 84-nucleotide deletion excises an AUG associated with a stronger Kozack sequence than the proline reading frame.¹³ Because the AUG for the polyproline frame is retained and competition is reduced, this potentially enhances the polyproline reading frame on *ASFMR1-TV2*.¹³ Relative expression of *ASFMR1-TV2*, nonspliced *ASFMR1*, total *ASFMR1*, and total *FMR1* mRNA were quantified using the VIIA 7 real-time quantitative PCR system (Applied Biosystems). The results were calculated using a "Delta-Delta CT" ($\Delta\Delta CT$) algorithm (first Delta CT [ΔCT] = [target expression] – [Gus-B expression]; second Delta CT [$\Delta\Delta CT$] = [ΔCT] – normal control [NC]; fold increase of target vs NC = $2^{-(\text{second delta CT})} = 2^{-\Delta\Delta CT}$; fold increase = $2^{-\Delta\Delta CT}$) to compare relative expressions of target mRNA to the expression of the GUS-B control mRNA

and then to the expression of an NC sample (primers and probes in table e-1, links.lww.com/NXG/AS8).

Statistical analysis

Analysis of variance (ANOVA) or the Kruskal-Wallis test (where appropriate) was performed to compare the levels of *ASFMR1-TV2*, nonspliced *ASFMR1*, total *ASFMR1*, and total *FMR1* among NCs, PMCs without FXTAS (PMC), and PMCs with FXTAS (FXTAS). The sample size calculation was based on ANOVA comparing NCs, PMC without FXTAS, and FXTAS. Pairwise comparisons were conducted with Bonferroni adjustment for multiple comparisons. The FXTAS-RS scores were compared among different clinical phenotypes with ANOVA or the Kruskal-Wallis test. Receiver operating characteristic (ROC) curve analysis was performed to evaluate the performance of *ASFMR1-TV2* to discriminate patients with FXTAS from others (NC and PMC). The following guide was used for interpretation: excellent 0.9–1; good 0.8–0.9; fair 0.7–0.8; poor 0.6–0.7; and fail 0.5–0.6. The Spearman correlation was used to examine the relationship between the CGG repeat size and the ratio of the splice variant and total *ASFMR1* mRNA (table e-2, links.lww.com/NXG/AS8). Least absolute shrinkage and selection operator (LASSO) logistic regression was performed to examine the association of the splice variant, CGG repeat size, AGG interruptions, nonspliced *ASFMR1*, total *ASFMR1*, *FMR1*, and family history of FXS with FXTAS adjusted for age. LASSO regression was used in this study to select the strongest predictors. AR was included in the regression for woman. Data for men and women were analyzed separately because the penetrance of FXTAS is much higher in men than in women. Missing data were excluded from the analysis.

Results

One hundred thirty-five participants including 56 men (mean, SD years; 65.2 ± 12.1 years) and 79 women (55.2 ± 15.5 years) were recruited.

Men

Men comprised 66% of those participants who met the criteria for FXTAS, 21% of PMCs without FXTAS and 45% of NCs. Twenty-six participants who met the criteria for FXTAS, with a mean age of 71 ± 8.4 years, had a median CGG repeat length of 91.5 (interquartile range [IQR] 29) (table 1). Twelve PMC participants without FXTAS, with a mean age of 61 ± 11.8 years, had a median CGG repeat length of 78.5 (IQR 21). Eighteen NCs had a mean age of 59.7 ± 13.6 years and a median CGG repeat length of 30 (IQR 10). Men with FXTAS were older than PMC and NC, and the CGG repeat size was higher in men with FXTAS than in PMC men and NC men. Nineteen (73.1%) men with FXTAS, 3 PMCs (25%), and 2 (11.8%) NC men had complete loss of AGG interruptions. FXTAS-RS scores were higher in FXTAS participants (median [IQR]: 56.5 [42]) than in the other 2

Table 1 Demographics in men (n = 56)

	NC, N = 18	PMC, N = 12	FXTAS, N = 26	p Value
Age, mean (SD)	59.72 (13.64)	61 (11.81)	70.96 (8.38)	0.003
CGG, median (IQR)	30 (10)	78.5 (21)	91.5 (29)	0.15 ^a
AGG, n (%)				0.0001
0	2 (11.76) ¹	3 (25)	19 (73.08)	
≥1	15 (88.24)	9 (75)	7 (26.92)	
FXTAS-RS, median (IQR)	5 (8)	7 (9.5)	56.5 (42)	<0.0001
Neuropathy, median (IQR)	0 (0) ⁷	4 (5) ⁸	6 (5) ⁹	0.002
Family history, n (%)	8 (44.44)	11 (91.67)	22 (84.62)	0.003
FXTAS diagnosis (%)				
Possible			2 (7.69)	
Probable			10 (38.46)	
Definite			14 (53.85)	

Abbreviations: FXTAS = fragile X-associated tremor/ataxia syndrome; FXTAS-RS = fragile X-associated tremor/ataxia syndrome rating scale; IQR = interquartile range; NC = normal control; PMC = premutation carrier.

Superscripts indicate the number of missing values.

^a Comparison between premutation and FXTAS.

groups ($p < 0.0001$). The total (modified) neuropathy score was 6 (IQR 5) in those with FXTAS, 4 (IQR 5) in PMCs and 0 (IQR 0) in NCs. Eighty-four percent ($p = 0.003$) of men with FXTAS had a family history of FXS. Participants who met the diagnostic criteria for FXTAS included 2 with possible, 10 with probable, and 14 with definite FXTAS. In men, there was a difference in all molecular variables ($p < 0.01$) among the 3 groups (table 2). Pairwise comparisons of the molecular results were only significant when comparing normal vs FXTAS and normal vs PMC, but not when comparing PMC vs FXTAS. ROC curve analysis indicated that *ASFMR1-TV2* has a good discriminating power of FXTAS in men (figure 1, A-F). The area under the curve (AUC) was 0.80 (95% CI = 0.68, 0.92). LASSO logistic regression analysis with the adjustment of age showed that loss of AGG (coefficient = -0.55) has strongest relationship with FXTAS, followed by a larger CGG repeat size (coefficient = 0.47) and an elevated *ASFMR1-TV2* level (coefficient = 0.21).

Coefficients for other independent variables including family history of FXS, nonspliced *ASFMR1*, total *ASFMR1*, and *FMR1* were shrunken to zero by the LASSO model, which indicates that these variables did not have a strong relationship with FXTAS.

Women


Thirteen participants who met the criteria for FXTAS, with a mean age of 69.8 ± 11 years, had an average CGG repeat length of 92.2 ± 15.1 (table 3). Forty-four PMCs without FXTAS, with a mean age of 50.3 ± 14.4 years, had an average CGG repeat length of 89.5 ± 19.6 . Twenty-two NCs, with a mean age of 56.4 ± 14.5 years, had an average CGG repeat length of 31.6 ± 4.7 . Women with FXTAS were older than PMC and NC, and the CGG repeat size was higher in women with FXTAS than in PMC women and NC women. Thirteen (100%) women with FXTAS, 32 PMCs (71%), and 2 (9%) healthy women had complete loss of AGG

Table 2 Molecular data in men (n = 56)

	NC, n = 18	PMC, n = 12	FXTAS, n = 26	p Value
<i>ASFMR1-TV2</i> , median	1.11 (1.42)	11.35 (20.22)	13.63 (15.74)	<0.0001
Nonspliced <i>ASFMR1</i> , mean	0.99 (0.59)	2.17 (1.54)	1.77 (1.22)	0.02
Total <i>ASFMR1</i> , mean	1.04 (0.78)	4.23 (3.26)	4.27 (2.91)	0.0002
Total <i>FMR1</i> , median	0.94 (0.44)	1.43 (0.46) ¹	1.21 (0.51) ¹	0.02

Abbreviations: *ASFMR1-TV2* = antisense fragile-X mental retardation transcript variant 2; *FMR1* = fragile X mental retardation 1; FXTAS = fragile X-associated tremor/ataxia syndrome; NC = normal control; PMC = premutation carrier.

Figure 1 ROC curve of *ASFMR1*-TV2



(A and B) ROC analysis indicated that *ASFMR1*-TV2 has a good discriminating power of FXTAS when comparing FXTAS with PMC and normal in men (A: AUC = 0.80; 95% CI = 0.68, 0.92) and women (B: AUC = 0.80; 95% CI = 0.67, 0.9). (C and D) ROC analysis indicated that *ASFMR1*-TV2 did not have a good discriminating power when comparing FXTAS with PMC in both men (C) and women (D) (men: AUC = 0.53, 95% CI = 0.32, 0.74; women: AUC = 0.69, 95% CI = 0.53, 0.85). (E and F) ROC analysis indicated that *ASFMR1*-TV2 has a good discriminating power when comparing PMC without FXTAS with normal in both men (E) and women (F) (men: AUC = 0.95, 95% CI = 0.86, 1; women: AUC = 0.92, 95% CI = 0.85, 0.99). (A–F) The green line is the diagonal reference line, and the blue line is the empirical ROC curve comparing FXTAS with NCs and premutation carriers (A and B), FXTAS with PMC (C and D), and PMC without FXTAS with controls (E and F) (The closer the ROC curve gets to the reference line, the worse the test. A large value of the AUC indicates a good test. An area of 1 means a perfect test, and 0.5 represents a worthless test). AUC = area under the curve; *ASFMR1*-TV2 = antisense fragile-X mental retardation-transcript variant 2; CI = confidence interval; FXTAS = fragile X-associated tremor/ataxia syndrome; NC = normal control; PMC = premutation carrier; ROC = receiver operating characteristic.

interruptions. FXTAS-RS scores were higher in FXTAS participants (median [IQR]: 30 [26]) than in the other 2 groups ($p < 0.0001$). The total (modified) Neuropathy Scale score was 2 (IQR = 2) in those with FXTAS, 1 (IQR = 3) in PMC and 0 (IQR = 1) in NC women ($p = 0.01$). More than 90% of women with FXTAS and PMCs had a family history of FXS. Participants who met the diagnostic criteria for

FXTAS included 3 with possible, 4 with probable, and 6 with definite FXTAS. In women, *ASFMR1*-TV2 levels, total *ASFMR1* levels, and *FMRI* levels were different among the groups (table 4). There were no differences in the non-spliced *ASFMR1* levels among the groups. In women, pairwise comparisons of the molecular results were significant when comparing between NC vs FXTAS and NC vs PMC,

Table 3 Demographics in women (n = 79)

	NC, n = 22	PMC, n = 44	FXTAS, n = 13	p Value
Age, mean (SD)	56.36 (14.52)	50.29 (14.40)	69.77 (11.01)	0.0003
CGG, mean (SD)	31.59 (4.68)	89.52 (19.64)	92.15 (15.07)	0.66 ^a
AR, mean (SD)	NA	0.50 ¹ (0.23)	0.46 (0.31)	0.87 ^a
AGG, n (%)				<0.0001
0	2 (9.09)	32 (72.73)	13 (100)	
≥1	20 (90.91)	12 (27.27)	0	
FXTAS-RS, median (IQR)	3 (4)	7 (8)	30 (26)	<0.0001
Neuropathy, median (IQR)	0 (1)	1 (3)	2 (2)	0.01
Family history, n (%)	5 (22.73)	40 (90.91)	12 (92.31)	<0.0001
FXTAS diagnosis (%)				
Possible			3 (23.08)	
Probable			4 (30.77)	
Definite			6 (46.15)	

Abbreviations: AR = activation ratio; FXTAS = fragile X-associated tremor/ataxia syndrome; FXTAS-RS = fragile X-associated tremor/ataxia syndrome rating scale; IQR = interquartile range; NC = normal control; PMC = premutation carrier. Superscripts indicate the number of missing values.

^a Comparison between premutation and FXTAS.

but were not when comparing PMC vs FXTAS. *ASPMR1-TV2* showed a good discriminating power of FXTAS from the ROC curve analysis (figure 2, AUC = 0.8, 95% CI = 0.67, 0.9). LASSO logistic regression, adjusting for age, showed that the loss of AGG (coefficient = -0.5555) has the strongest relationship with FXTAS, followed by *ASPMR1-TV2* (coefficient = 0.4747) and total *ASPMR1* (coefficient = 0.2121). Other variables including family history of FXS, nonspliced *ASPMR1*, AR, CGG repeat size, and *FMR1* did not show a strong relationship with FXTAS from the LASSO regression model.

In men, we found a negative correlation between the CGG repeat size and the ratio of the splice variant and total *ASPMR1* mRNA among the NCs (correlation coefficient = -0.62, $p = 0.04$), but a positive correlation among the PMCs (correlation coefficient = 0.81, $p < 0.0001$). In women, there

was also a negative correlation among the controls (correlation coefficient = -0.74, $p = 0.02$), but no correlation was found among the PMCs (CGG was adjusted for AR) (correlation coefficient = 0.08, $p = 0.70$).

Discussion


The main purpose of this study was to identify molecular markers to help predict which PMC will get FXTAS because even for male carriers, not all PMC will get the disease.¹⁴ The exact clinical impact of *ASPMR1* splice variants is unclear; however, it has been postulated that in addition to *FMR1*, elevated levels of the expanded GCC repeat containing the *ASPMR1* transcript are observed in PMCs and may contribute to the variable phenotypes associated with the CGG repeat expansion.¹⁴

Table 4 Molecular results in women (n = 79)

	NC, n = 22	PMC, n = 44	FXTAS, n = 13	p Value
<i>ASPMR1-TV2</i> , median	0.91 (0.67)	6.92 (10.29)	12.52 (29.37)	<0.0001
Nonspliced <i>ASPMR1</i> , mean	0.78 (0.41)	0.93 (0.69)	1.3 (0.96)	0.1
Total <i>ASPMR1</i> , mean	0.93 (0.48)	2.42 (1.87)	3.52 (2.62)	0.0002
Total <i>FMR1</i> , median	0.89 (0.32)	1.05 (0.48)	1.22 (0.43)	0.01

Abbreviations: *ASPMR1-TV2* = antisense fragile-X mental retardation transcript variant 2; *FMR1* = fragile X mental retardation 1; FXTAS = fragile X-associated tremor/ataxia syndrome; NC = normal control; PMC = premutation carrier.

Figure 2 Antisense FMR transcript variant 2



Defining characteristic of *ASFMR1-TV2*. There is an 84-nt deletion at near the beginning of the *ASFMR1* ORF that has an enhanced expression compared with a healthy control. We designed strand specific primers that target the presence and absence of the 84-nt sequence to determine the fold increase relative to *GUS* expression for *ASFMR1-TV2* spliced and nonspliced assays. *ASFMR1-TV2* = antisense fragile-X mental retardation transcript variant 2; FMR = fragile X mental retardation; nt = nucleotide; ORF = open reading frame.

This is the first study to evaluate *ASFMR1* splice variant levels in PMCs and FXTAS patients. The reason this study concentrated on *ASFMR1-TV2* (figure 2) was because small unpublished studies suggested that this splice variant of the *ASFMR1* transcript might be associated with neurologic symptoms in FXTAS men. This study reports elevated levels of *ASFMR1-TV2* in FXTAS and a loss of AGG interruptions in men with FXTAS. In addition, this study also showed that the FXTAS-RS and Neuropathy Scale scores were higher in patients with FXTAS compared with PMCs and NCs. We were not able to distinguish between PMCs and FXTAS with pairwise comparisons of *ASFMR1-TV2*, nonspliced *ASFMR1*, total *ASFMR1*, or total *FMR1*.

Although we found that *ASFMR1-TV2* and lack of AGG were more common in FXTAS, the power to discriminate between PMCs and FXTAS was not high enough to use alone diagnostically or in a predictive manner. This may be because our sample sizes were too small overall. Some of the groups, such as the PMC men without FXTAS and the FXTAS women group did not have many participants. Because the disease is highly penetrant in men, it was difficult to recruit PMC men without FXTAS. Women with FXTAS were also difficult to recruit because there is a lower prevalence in women. The published criteria for FXTAS^{3,4} were followed, and because the women have milder features than men with FXTAS, many of them needed either the middle cerebellar peduncle sign or brainstem white matter lesions to qualify for probable FXTAS. Thus, the strict application of the diagnostic

criteria leads to the high number having the MCP sign rather than our population of FXTAS women being different from the populations that have been published in the past. Alternatively, the patterns of relationships found may indicate that regardless of the sample size, elevated RNA products from the repeat-containing gene are a phenomenon more related to just the presence of a premutation, rather than reflecting a threshold of abnormal RNAs at which symptoms occur. RAN translation has been described in FXTAS as a potential additional mechanism of toxicity independent of the presumed RNA toxicity. Measurement of RAN translation products could be more closely correlated with amount and on the time of the onset of symptoms. Additional studies may be warranted to combine molecular and other biomarkers, such as examination, imaging, or laboratory markers, to increase the positive predictive value and the clinical utility.

Both AUG-initiated and repeat-associated non-AUG (RAN) translation have been shown to occur from *ASFMR1* CCG repeats.^{2,8} Associated transcripts are capable of forming strong secondary structures in vitro, either RNA hairpins or G-quadruplexes, and are important to the process underlying RAN translation. It is yet unknown if secondary structures linked to *ASFMR1-TV2* may also be linked to RAN translation and whether toxicity occurs from the expression of proteins generated from translation of this variant. Pilot work from our research group suggests an association between the lack of AGG interspersions in *FMR1* PMCs and higher rates of neurologic signs, which was confirmed in this study. It is

recognized that the results regarding the lack of AGG interruptions may represent instead a relationship with the longest pure CGG repeat size. The loss of AGG interruptions is seen in PMCs at a much higher rate than NCs because it makes these repeats more unstable and hence more likely to expand meiotically and (presumably) somatically in tissues. Similarly, one would presume that PMCs with fewer AGG interruptions would have larger pure repeats. Then, these larger pure repeats are associated with enhanced transcription of the *FMRI* and *ASFMRI* loci because of changes in local chromatin.^{29,30} Additional studies are needed to determine which factor is responsible for the association.

It is important to find molecular markers that can serve as screening tools and increase diagnostic certainty, not only to help families of patients at risk of disorder but also to enable drug discovery, so timely treatment can be implemented. Both PMCs and FXTAS participants had elevated levels of *ASFMRI* splice variant 2, and loss of AGG interruptions was associated with FXTAS in men. These markers may have utility if combined with other predictive factors to better predict who will get FXTAS. Additional work also needs to be done to determine the contribution to the mechanism of CCG-mediated toxicity in PMCs.

Author contributions

P. Vittal: conception of the study, subject recruitment and neurologic evaluation, skin biopsy, and organizing and writing of the first draft of the manuscript. S. Pandya and K. Sharp: performance of RNA assays. J. Jackson: performance of genotyping and AGG analysis. L. Zhou: performance of genotyping and AGG analysis, fibroblast cultures, and supervision of laboratory assays. B. Ouyang: statistical analysis. E. Berry-Kravis: conception of the study, review and critique of the manuscript, and oversight of laboratory studies. D.A. Hall: conception of the study, subject recruitment, and review and critique of the manuscript.

Acknowledgment

The authors thank the American Brain Foundation and the National Ataxia Foundation for their support of the Clinical Research Training Fellowship grant in Ataxia and Tracy Waliczek and Evelyn Perez for their assistance during skin biopsy and blood sample collection. They thank Integrated DNA Technologies (Coralville, IA) for custom-designed primers and probes. They also thank the participants of this study.

Study funding

Clinical Research Training Fellowship grant in Ataxia sponsored by the American Brain Foundation and the National Ataxia Foundation.

Disclosure

P. Vittal has received funding for travel and/or speaker honoraria and served on speakers' bureaus of Teva Pharmaceutical Industries Ltd and Adamas Pharmaceuticals Inc. and has

received research support from the American Brain Foundation and the National Ataxia Foundation. S. Pandya and K. Sharp report no disclosures. E. Berry-Kravis serves or has served on the scientific advisory boards of Vteese, BioMarin, Marinus, Fulcrum, Cydan, GW, and Ovid; holds a patent regarding a method for assay of CCHS-causing polyalanine repeat expansion for diagnosis; receives or has received publishing royalties for The Fragile X Tremor Ataxia Syndrome (FXTAS); consults or has consulted for Fulcrum and Zynerba; and has received research support from Vtesse, Ovid, Neuren, NIH/NICHHD, NIH/NCATS, CDC, NIH/NINDS, NIH/NIMH, and the John Merck Fund. L. Zhou, B. Ouyang, and J. Jackson report no disclosures. D.A. Hall served or has served on the scientific advisory board of the National Fragile X Foundation; serves or has served on the editorial board of *Neurology Today*; consults or has consulted for the NIH; and has received research support from Pfizer, Neurocrine, AbbVie, NIH, Shapiro Foundation, Anti-Aging Foundation, and Parkinson Foundation. Full disclosure form information provided by the authors is available with the full text of this article at Neurology.org/NG.

Received August 14, 2017. Accepted in final form April 11, 2018.

References

1. Hagerman RJ, Leehey M, Heinrichs W, et al. Intention tremor, parkinsonism, and generalized brain atrophy in male carriers of fragile X. *Neurology* 2001;57:127–130.
2. Hall DA, O'Keefe JA. Fragile x-associated tremor ataxia syndrome: the expanding clinical picture, pathophysiology, epidemiology, and update on treatment. *Tremor Other Hyperkinet Mov (N Y)* 2012;2.
3. Jacquemont S, Hagerman RJ, Leehey M, et al. Fragile X premutation tremor/ataxia syndrome: molecular, clinical, and neuroimaging correlates. *Am J Hum Genet* 2003;72:869–878.
4. Brunberg JA, Jacquemont S, Hagerman RJ, et al. Fragile X premutation carriers: characteristic MR imaging findings of adult male patients with progressive cerebellar and cognitive dysfunction. *Am J Neuroradiol* 2002;23:1757–1766.
5. Pieretti M, Zhang FP, Fu YH, et al. Absence of expression of the *FMR-1* gene in fragile X syndrome. *Cell* 1991;66:817–822.
6. Tassone F, Long KP, Tong TH, et al. *FMR1* CGG allele size and prevalence ascertained through newborn screening in the United States. *Genome Med* 2012;4:100.
7. Tassone F, Bellina A, Carosi C, et al. Elevated *FMR1* mRNA in premutation carriers is due to increased transcription. *RNA* 2007;13:555–562.
8. Todd PK, Oh SY, Krans A, et al. CGG repeat-associated translation mediates neurodegeneration in fragile X tremor ataxia syndrome. *Neuron* 2013;78:440–455.
9. Robin G, López JR, Espinal GM, et al. Calcium dysregulation and Cdk5-ATM pathway involved in a mouse model of fragile X-associated tremor/ataxia syndrome. *Hum Mol Genet* 2017;26:2649–2666.
10. Leehey MA, Berry-Kravis E, Goetz CG, et al. *FMR1* CGG repeat length predicts motor dysfunction in FXTAS. *Neurology* 2008;70:1397–1402.
11. Berry-Kravis E, Potanos K, Weinberg D, et al. Penetrance of fragile X-associated tremor/ataxia syndrome (FXTAS) in two sisters related to X-inactivation pattern. *Ann Neurol* 2005;57:144–147.
12. Hall DA, Robertson-Dick EE, O'Keefe JA, et al. Repeat size and X-inactivation in the clinical phenotype of fragile X premutation carrier sisters: a familial case series. *Neurol Genet* 2016;2:e45. doi: 10.1212/NXG.0000000000000045.
13. Ladd PD, Smith LE, Rabaia NA, et al. An antisense transcript spanning the CGG repeat region of *FMR1* is upregulated in premutation carriers but silenced in full mutation individuals. *Hum Mol Genet* 2007;16:3174–3187.
14. Yrigollen CM, Durbin-Johnson B, Gane L, et al. AGG interruptions within the maternal *FMR1* gene reduce the risk of offspring with fragile X syndrome. *Genet Med* 2012;14:729–736.
15. Hall D, Todorova-Koteva K, Pandya S, et al. Neurological and endocrine phenotypes of fragile X carrier women. *Clin Genet* 2016;89:60–67.
16. Berry-Kravis E, Goetz CG, Leehey MA, et al. Neuropathic features in fragile X premutation carriers. *Am J Med Genet A* 2007;143A:19–26.
17. Jacquemont S, Hagerman RJ, Leehey MA, et al. Penetrance of the fragile X-associated tremor/ataxia syndrome in a premutation carrier population. *JAMA* 2004;291:460–469.
18. Leehey M, Goetz C, Berry-Kravis E. Development of the FXTAS rating scale for quantitative motor analysis. Presented at the 10th International Congress of Parkinson's Disease and Movement Disorders; October 30, 2006; Kyoto, Japan.

19. Cornblath DR, Chaudhry V, Carter K, et al. Total neuropathy score: validation and reliability study. *Neurology* 1999;53:1660–1664.
20. Fahn S, Tolosa E, Marin C. Clinical rating scale for tremor. In: Jankovic J, Tolosa E, editors. *Parkinson's Disease and Movement Disorders*. Baltimore, Munich: Urban & Schwarzenberg; 1987:225–234.
21. Trouillas P, Takayanagi T, Hallett M, et al. International cooperative ataxia rating scale for pharmacological assessment of the cerebellar syndrome: the Ataxia Neuropharmacology Committee of the World Federation of Neurology. *J Neurol Sci* 1997;145:205–211.
22. Fahn S, Elton R; members of the UPDRS Development Committee. Unified Parkinson's disease rating scale. In: Fahn S, Marsden C, Calne D, Goldstein M, editors. *Recent Development in Parkinson's Disease*. Florham Park, NJ: Macmillan Health Care Information; 1987:153–164.
23. Siesling S, Zwinderman AH, van Vugt JP, et al. A shortened version of the motor section of the Unified Huntington's Disease Rating Scale. *Mov Disord* 1997;12:229–234.
24. Chen L, Hadd A, Sah S, et al. An information-rich CGG repeat primed PCR that detects the full range of fragile X expanded alleles and minimizes the need for southern blot analysis. *J Mol Diagn* 2010;12:589–600.
25. Filipovic-Sadic S, Sah S, Chen L, et al. A novel FMR1 PCR method for the routine detection of low abundance expanded alleles and full mutations in fragile X syndrome. *Clin Chem* 2010;56:399–408.
26. Godler DE, Tassone F, Loesch DZ, et al. Methylation of novel markers of fragile X alleles is inversely correlated with FMRP expression and FMR1 activation ratio. *Hum Mol Genet* 2010;19:1618–1632.
27. Annemieke JMHV, Pieretti M, Sutcliffe JS, et al. Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* 1991;65:905–914.
28. Krans A, Kearse M, Todd PK. Repeat-associated non-AUG translation from antisense CCG repeats in fragile X tremor/ataxia syndrome. *Ann Neurol* 2016;80:871–881.
29. Todd PK, Oh SY, Krans A, et al. Histone deacetylases suppress CGG repeat-induced neurodegeneration via transcriptional silencing in models of fragile X tremor ataxia syndrome. *PLoS Genet* 2010;6:e1001240.
30. Hoem G, Raske CR, Garcia-Arocena D, et al. CGG-repeat length threshold for FMR1 RNA pathogenesis in a cellular model for FXTAS. *Hum Mol Genet* 2011;20:2161–2170.

Neurology[®] Genetics

ASPMRI splice variant: A predictor of fragile X-associated tremor/ataxia syndrome

Padmaja Vittal, Shrikant Pandya, Kevin Sharp, et al.

Neurol Genet 2018;4;

DOI 10.1212/NXG.0000000000000246

This information is current as of July 30, 2018

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/4/4/e246.full.html
References	This article cites 26 articles, 4 of which you can access for free at: http://ng.neurology.org/content/4/4/e246.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Gait disorders/ataxia http://ng.neurology.org/cgi/collection/gait_disorders_ataxia Tremor http://ng.neurology.org/cgi/collection/tremor
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2018 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.

