Are migraineurs naturally born “well-hearted”?

OPEN

Anne Ducros, MD, PhD

Correspondence to
Prof. Ducros;
a-ducros@chu-montpellier.fr

Neurol Genet
2015;1:e8; doi: 10.1212/
NXG.0000000000000008

Migraine is a common episodic brain disorder characterized by recurrent attacks of disabling headache.¹ In one-third of patients, headache is preceded or accompanied by aura symptoms, which are transient and fully reversible hemispheric neurologic symptoms. In the absence of any objective marker, the diagnosis of migraine is clinical and based on the criteria proposed by the International Headache Society, which distinguish migraine without aura (patients who have never had an aura) and migraine with aura (patients who have had at least 2 auras, irrespective of the number of attacks without aura they have had).¹

Migraine affects about 15% of the occidental population and predominates in women. Family and twin studies have shown that migraine is polygenic, with an overall heritability nearing 50%.²

Migraine is viewed as a benign condition, probably because the migraineurs who seek care are mostly young or middle-aged adults and because attacks tend to improve and eventually disappear with aging. However, accumulating data show that migraine is an independent risk factor for vascular disorders. Clinic- and population-based studies have shown that migraine with aura (and not migraine without aura) doubles the risk of ischemic stroke, especially in women younger than 45 years who are free of any other vascular risk factors (with the exception of smoking and use of oral contraceptives, which further increase the risk).^{3,4} Cortical spreading depression, the surrogate of migraine aura, has been proposed as the main link between migraine and ischemic stroke through an increased vulnerability to cerebral ischemia.⁵ Several studies and a recent meta-analysis have also shown that migraine, especially migraine with aura, is associated with an increased risk of coronary artery diseases (CADs), including angina, myocardial infarction, and coronary revascularization procedures.⁶ The mechanisms linking migraine to CAD are totally unknown.

In this issue of *Neurology® Genetics*, Winsvold et al.⁷ studied shared genetic susceptibility underlying

migraine and CAD. Their main hypothesis was that shared genetic variations might explain part of the observed epidemiologic link between migraine and CAD. The authors analyzed 2 large existing genome-wide association studies (GWAS) for migraine and CAD. Using 4 separate methods, they found that some genetic loci are indeed associated with both migraine and CAD. However, and probably contrary to expectations, they found no overlap of genetic risk loci between migraine with aura and CAD. Moreover, the genetic variations that overlap between migraine without aura and CAD are protective.⁷ In other words, patients with migraine without aura seem to have a lower load of genetic factors increasing the risk of CAD, and patients with migraine with aura do not have more risk loci increasing the risk of CAD than controls.

We now need to understand why migraineurs who are born with a protective or neutral genetic risk profile for CAD end up with an increased risk of coronary events. Other genetic factors could play a role. GWAS are efficient at identifying common genetic variants that each have a small functional effect. The most recent meta-analysis of GWAS has identified 42 variants associated with migraine, of which 7 are specific for migraine without aura and none is specific for migraine with aura.⁸ One hypothesis is that the mode of inheritance of migraine with aura differs from the complex polygenic inheritance of migraine without aura and involves rare variants that each exert more important functional effects. Furthermore, it has been suggested that migraine with aura could be associated not only with an increased neuronal excitability and vulnerability to cerebral ischemia because of the consequences of cortical spreading depression but also with an increased cardiac and cellular vulnerability.⁹ Some rare genetic variants, which cannot be captured by the GWAS approach, could be implicated in both migraine with aura and CAD. Second, nongenetic factors could play a role. Migraine has been associated with adverse life style,

See article

From the Department of Neurology, Neuroscience Head and Neck Clinic, Montpellier University Hospital, and the University of Montpellier, Montpellier, France.

Funding information and disclosures are provided at the end of the editorial. Go to Neurology.org/ng for full disclosure forms. The Article Processing Charge for this editorial was waived at the discretion of the Editor.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

obesity, avoidance of physical activity, smoking, unfavorable cholesterol profiles, and depression, which all increase the risk of CAD.⁴ Hypercoagulable states and endothelial dysfunction, both of which have been associated with migraine, could also play a role.⁴ It is important to emphasize that the use of triptans and ergots to abort migraine headache cannot explain the epidemiologic link between migraine and CAD.¹⁰ These vasoconstrictive migraine medications are contraindicated in patients who have had a stroke or a CAD or who have uncontrolled hypertension.

Practically, physicians should inform patients affected by migraine with aura of the increased risk of stroke and CAD, detect and treat conventional modifiable vascular risk factors, and especially encourage smoking cessation and regular physical activity.

While more data are needed to understand the molecular mechanisms underlying the association between migraine and CAD, this study shows that common genetic variants are not major players in this association.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

A. Ducros serves as associate editor of *Cephalalgia* and *Headache* and as a member of the editorial advisory board of *The Journal of Headache and Pain*; received travel expenses and honoraria for an advisory board from Amgen; received travel expenses and/or honoraria for lectures or educational activities not funded by industry; and receives institutional support from the French government (Contrat d'Initiation à la Recherche Clinique, Programme Hospitalier de Recherche Clinique) for clinical research. Go to Neurology.org/ng for full disclosure forms.

REFERENCES

1. Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia* 2013;33:629–808.
2. Ferrari MD, Klever RR, Terwindt GM, Ayata C, van den Maagdenberg AM. Migraine pathophysiology: lessons from mouse models and human genetics. *Lancet Neurol* 2015;14:65–80.
3. Kurth T, Chabriat H, Bousser MG. Migraine and stroke: a complex association with clinical implications. *Lancet Neurol* 2012;11:92–100.
4. Mawet J, Kurth T, Ayata C. Migraine and stroke: in search of shared mechanisms. *Cephalalgia* 2015;35:165–181.
5. Eikermann-Haerter K. Spreading depolarization may link migraine and stroke. *Headache* 2014;54:1146–1157.
6. Sacco S, Ornello R, Ripa P, et al. Migraine and risk of ischemic heart disease: a systematic review and meta-analysis of observational studies. *Eur J Neurol* 2015;22:1001–1011.
7. Winsvold BS, Nelson CP, Malik R, et al; for the CARDIoGRAM Consortium and the International Headache Genetics Consortium. Genetic analysis for a shared biological basis between migraine and coronary artery disease. *Neurol Genet* 2015;1:e10. doi: 10.1212/NXG.000000000000010.
8. Anttila V, Gormley P, Muona M, Neale BM, Palotie A; International Headache Genetics Consortium. Genome-wide meta-analysis of migraine in 59,000 cases and 316,000 controls reveals 29 new loci, increasing the total number of known risk loci to 42. *Cephalalgia* 2015;35(6 suppl):1–296, OR10.
9. Ripa P, Ornello R, Pistoia F, Carolei A, Sacco S. Spreading depolarization may link migraine, stroke, and other cardiovascular disease. *Headache* 2015;55:180–182.
10. Roberto G, Raschi E, Piccinini C, et al. Adverse cardiovascular events associated with triptans and ergotamines for treatment of migraine: systematic review of observational studies. *Cephalalgia* 2015;35:118–131.

Neurology[®] Genetics

Are migraineurs naturally born "well-hearted"?

Anne Ducros

Neurol Genet 2015;1;

DOI 10.1212/NXG.0000000000000008

This information is current as of July 2, 2015

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/1/1/e8.full.html
References	This article cites 10 articles, 1 of which you can access for free at: http://ng.neurology.org/content/1/1/e8.full.html#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://ng.neurology.org/content/1/1/e8.full.html#otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Cerebrovascular disease/Stroke http://ng.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke Association studies in genetics http://ng.neurology.org/cgi/collection/association_studies_in_genetics Cardiac http://ng.neurology.org/cgi/collection/cardiac Migraine http://ng.neurology.org/cgi/collection/migraine Risk factors in epidemiology http://ng.neurology.org/cgi/collection/risk_factors_in_epidemiology
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.