Qi Niu, MD
Xingxia Wang, MD
Mingchao Shi, MD
Qingwen Jin, MD

Neurol Genet
2015;1:e20; doi: 10.1212/
NXG.000000000000017

A NOVEL *DYNC1H1* MUTATION CAUSING SPINAL MUSCULAR ATROPHY WITH LOWER EXTREMITY PREDOMINANCE

OPEN

Recent studies have identified mutations in the dynein heavy chain gene (*DYNC1H1*), which lead to 2 closely related human motor neuropathies: a dominant spinal muscular atrophy with lower extremity predominance (SMALED) and axonal Charcot-Marie-Tooth (CMT) disease.^{1,2} We describe the identification of a novel mutation (p.G807S) in *DYNC1H1* as the cause of SMALED.

Case report. This study received approval from the People's Hospital of Jiangsu Province Human Studies Committee Institutional Review Board for experiments using human subjects. We obtained written informed consent from all subjects (or their guardians) participating in the study. The affected boy was born after 41 weeks of gestation. Pregnancy was uneventful, and birth weight was 3,080 g. His initial development was normal, and head control was recognized at 3–4 months. Late infantile motor development was mildly delayed, and he could walk unassisted at 2 years. An unstable gait persisted thereafter, and he was referred to us at 3 years and 6 months of age for evaluation. On examination he was unable to run, climbed stairs on all fours, walked unsteadily, and fell frequently. In contrast, his arms showed age-appropriate strength and no sensory loss could be appreciated. No cognitive delay was present, and he required a normal educational plan in nursery school. The patient is currently 7 years old and attends a regular primary school.

EMG of the tibialis anterior and genioglossus muscles showed evidence of denervation and reinnervation, consistent with non-length-dependent motor neuron disease. Neither serum transaminase nor creatine kinase was elevated, and brain MRI revealed normal findings. Muscle CT demonstrated severe atrophy and lipid degeneration, predominantly in the bilateral quadriceps femoris muscle. However, the upper limbs and distal lower limbs were not affected. A muscle biopsy from the quadriceps femoris muscle demonstrated severe grouping atrophy of type 2 fibers, with a massive increase in the amount of

fibrous tissue and sparse enlarged type 1 fibers. Genomic analysis data showed a heterozygous mutation c.2419G>A in exon 8 of the *DYNC1H1* gene, causing a glycine to serine substitution (p.G807S).

Three members of the subject's family have been diagnosed with SMALED. In his mother and grandfather, disease development has steadily progressed and both show moderate proximal lower limb–dominant muscle weakness and atrophy. They can walk unassisted, and both show a waddling gait and positive Gower sign. No sensory disturbance or ataxia is noted, and they do not have any intellectual disability. The 3 patients showed no pathologic mutations in *SMN1*, *SMN2*, *UBA1*, or *VAPB*.

Discussion. The proband in this study presented with a quadriceps femoris muscle that was almost exclusively involved in the early course of the disease, and the proximal lower limb was predominantly involved throughout disease progression. The subject eventually developed early childhood onset of proximal leg weakness with muscle atrophy, and clinical findings support non-length-dependent motor neuron disease without sensory involvement. His presentation and clinical course were characteristic of SMALED. We found a missense mutation [p.G807S (c.2419G>A)] on exon 8 in *DYNC1H1* (figure, A), a mutation not previously been reported as causative for SMALED or CMT disease. The heterozygous missense mutation was confirmed in I-1, II-2, and III-2 (figure, A), and the position is highly conserved (figure, B). This missense mutation was not found in 187 control samples. Clinical features of affected individuals in this family are essentially consistent with those of SMALED, the hallmarks of which are early childhood onset of proximal leg weakness with muscle atrophy and non-length-dependent motor neuron disease without sensory involvement.

Thirteen missense mutations identified in human cases of CMT or SMALED are located in the same tail domain of *DYNC1H1*, which is involved in protein dimerization.^{3–5} To assess the consequence of mutations in the *DYNC1H1* tail domain, we examined the stability and function of dynein complexes purified from dermal fibroblasts derived from the proband and his parents. We observed that dynein

Figure *DYNC1H1* mutation in pedigree with dominant spinal muscular atrophy and effect on the dynein complex

(A) The pedigree of a family with a dominant spinal muscular atrophy with lower extremity predominance. The proband is marked by an arrow. (B) Alignment of the *DYNC1H1* protein showing high conservation of the mutated position in exon 8 of *DYNC1H1*. (C) Dynein from the proband and his parents' fibroblasts were incubated with microtubules in the presence of adenosine triphosphate. After centrifugation, binding of dynein was assessed by immunoblotting for intermediate chain using mouse monoclonal to dynein intermediate chain 1 antibody (ab6304, Abcam). Glyceraldehyde 3-phosphate dehydrogenase was used as internal control. Line 1: mother; line 2: father; line 3: proband.

complexes from cells containing the *DYNC1H1* G807S mutation (the proband and his mother) showed markedly decreased binding to microtubules in the presence of adenosine triphosphate, compared to complexes from the father (figure, C). Our data suggest that the G807S mutation may disrupt dynein complex formation in a dominant negative manner. We further sequenced the *DYNC1H1* whole genome and found a splicing variant in an intron involving the reciprocal third nucleotide before position 12685 (c.12685-3C>T) that coexisted with the G807S mutation in both the proband and his mother. This mutation could also affect the function of dynein complexes. Together, our clinical features illustrate a possible clinical-genetic phenotype of SMALED associated with newly identified *DYNC1H1* mutations. However, further data will be required to establish possible genotype-phenotype correlations.

From the Department of Neurology (Q.N., X.W., M.S., Q.J.), The People's Hospital of Jiangsu Province, Nanjing, China; and Department of Neurology (Q.J.), The First Affiliated Hospital, Nanjing Medical University, Nanjing, China.

Author contributions: Study concept and design: Q.N. and Q.J.; acquisition of data: Q.N. and X.W.; analysis and interpretation of data: Q.N. and M.S.; drafting of the manuscript: Q.N. and Q.J.; critical revision of the manuscript for intellectual content: Q.N. and Q.J.; technical and material support: M.S.; study supervision: Q.J.

Acknowledgment: The authors thank patients and controls for having contributed to this study.

Study funding: Funding for this study was provided by grants from the National Natural Science Foundation of China (#30870125) and the Jiangsu Province Natural Science Foundation (#SBK201402229).

Disclosure: Dr. Niu, Dr. Wang, Dr. Shi, and Dr. Jin report no disclosures. Go to Neurology.org/ng for full disclosure forms. The Article Processing Charge was paid by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Received March 18, 2015. Accepted in final form June 26, 2015.

Correspondence to Dr. Jin: qingwen_jin@126.com

1. Weedon MN, Hastings R, Caswell R, et al. Exome sequencing identifies a *DYNC1H1* mutation in a large pedigree with dominant axonal Charcot-Marie-Tooth disease. *Am J Hum Genet* 2011;89:308–312.

2. Harms MB, Ori-McKenney KM, Scoto M, et al. Mutations in the tail domain of DYNC1H1 cause dominant spinal muscular atrophy. *Neurology* 2012;78:1714–1720.
3. Eschbach J, Sinniger J, Bouitbir J, et al. Dynein mutations associated with hereditary motor neuropathies impair mitochondrial morphology and function with age. *Neurobiol Dis* 2013;58:220–230.
4. Punetha J, Monges S, Franchi ME, et al. Exome sequencing identifies DYNC1H1 variant associated with vertebral abnormality and spinal muscular atrophy with lower extremity predominance. *Pediatr Neurol* 2015;52:239–244.
5. Scoto M, Rossor AM, Harms MB, et al. Novel mutations expand the clinical spectrum of DYNC1H1-associated spinal muscular atrophy. *Neurology* 2015;84:668–679.

Neurology[®] Genetics

A novel *DYNC1H1* mutation causing spinal muscular atrophy with lower extremity predominance

Qi Niu, Xingxia Wang, Mingchao Shi, et al.

Neurol Genet 2015;1;

DOI 10.1212/NXG.0000000000000017

This information is current as of July 16, 2015

Updated Information & Services

including high resolution figures, can be found at:
<http://ng.neurology.org/content/1/2/e20.full.html>

References

This article cites 5 articles, 0 of which you can access for free at:
<http://ng.neurology.org/content/1/2/e20.full.html##ref-list-1>

Citations

This article has been cited by 8 HighWire-hosted articles:
<http://ng.neurology.org/content/1/2/e20.full.html##otherarticles>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Genetic linkage

http://ng.neurology.org/cgi/collection/genetic_linkage

Peripheral neuropathy

http://ng.neurology.org/cgi/collection/peripheral_neuropathy

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://ng.neurology.org/misc/about.xhtml#permissions>

Reprints

Information about ordering reprints can be found online:

<http://ng.neurology.org/misc/addir.xhtml#reprintsus>

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.