Aravindhan
Veerapandiyan, MBBS
Amit Chaudhari, PhD
Christin M. Traba, MD,
MPH
Xue Ming, MD, PhD

Neurol Genet
2016;2:e99; doi: 10.1212/
NXG.0000000000000099

NOVEL MUTATION IN MITOCHONDRIAL DNA IN 2 SIBLINGS WITH LEIGH SYNDROME

OPEN

Leigh syndrome is clinically and genetically heterogeneous, associated with mutations in mitochondrial and nuclear genes.¹ Diagnostic criteria include progressive disorder with motor and intellectual delay/regression; signs and symptoms of brainstem and/or basal ganglia disease; raised lactate concentration in blood and/or CSF; and one or more of the following: (1) characteristic features on neuroimaging (bilateral symmetrical hyperintensities in brainstem, basal ganglia, dentate nuclei, and optic nerves on T2-weighted MRI); (2) typical neuropathologic changes; and (3) typical neuropathology in a similarly affected sibling.² We describe 2 African American siblings who have a mutation in the mitochondrial *MT-TL2* gene and a clinical diagnosis of Leigh syndrome. The same mutation is also identified in their neurologically asymptomatic mother.

The brother who failed to progress on motor milestones since age 1 year presented with 4 months of progressive ataxia and weakness at 2 years of age. He was not able to pull to stand or walk without support. Growth parameters were below the third percentile. Examination revealed bilateral ptosis, left facial weakness, global hypotonia, mild weakness in all extremities, and severe truncal and limb ataxia. Serum and CSF lactate levels were 2.8 mmol/L (0.5–2.2) and 2.6 mmol/L (0.5–2.2), respectively. Brain MRI revealed T2 hyperintensities involving bilateral putamen, red nuclei, substantia nigra, thalami, and periaqueductal gray matter (figure). Mitochondrial DNA sequencing revealed a homoplasmic m.12311 T>C change on *MT-TL2* encoding tRNA Leu (CUN). Two months later, he required mechanical ventilation during sleep and subsequently died at the age of 3.5 years. His younger sister presented at 3 years of age with a 1-year history of motor regression, speech delay, and ataxia. She lost her ability to walk or stand unassisted, exhibiting truncal and limb ataxia, but was able to crawl and sit with support. She had limited 2-word phrases. Growth parameters were below the ninth percentile. Serum lactate level was 1.72 mmol/L (0.25–1.1). T2 hyperintensities were

seen in bilateral substantia nigra, thalami, and brainstem. She remains neurologically stable at the age of 4 years.

Association of this T-to-C substitution at nucleotide 12311 on *MT-TL2* with Leigh syndrome has not been clearly described. This mutation was previously described in a patient with chronic progressive external ophthalmoplegia.³ In addition, heteroplasmic *MT-TL2* mutations have been reported in patients with mitochondrial encephalomyopathy and myopathy.⁴ Affected site in our patients is conserved among species during evolution,^{5,6} and the gene itself is one of the most conserved mitochondrial *tRNA* genes among closely related species of hominoids.⁷ Mutations in mitochondrial *tRNA* genes were reported in Leigh syndrome.¹ We speculate that this mutation in the highly conserved region is associated with Leigh syndrome phenotypes in our patients. This is supported by the presence of the same *MT-TL2* mutation in their mother with bipolar disorder, indicating that this change may have variable penetrance or heterogeneous expressivity commonly seen with other mitochondrial disorders. We suggest that patients with clinical criteria of Leigh syndrome be screened for this mutation to confirm such an association.

From the Department of Neurology (A.V., A.C., S.M.), Division of Pediatric Neurology, and Department of Pediatrics (C.M.T.), Rutgers New Jersey Medical School, Newark.

Author contributions: Aravindhan Veerapandiyan: study concept, drafting, and revising the manuscript. Amit Chaudhari: drafting the manuscript. Christin M. Traba: revising the manuscript. Xue Ming: study concept and revising the manuscript.

Study funding: No targeted funding reported.

Disclosure: Dr. Veerapandiyan, Dr. Chaudhari, and Dr. Traba report no disclosures. Dr. Ming has served on the editorial boards of the *Journal of Pediatric Neurology* and the *Journal of Sleep Disorders and Therapy*; and holds a patent for Polyunsaturated fatty acids interactions and oxidative stress disorders. Go to Neurology.org/ng for full disclosure forms. The Article Processing Charge was paid by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Received April 28, 2016. Accepted in final form July 13, 2016.

Correspondence to Dr. Ming: mingxu@njms.rutgers.edu

Axial T2 image showing hyperintense lesions involving periaqueductal gray matter and cerebellar hemispheres in our first patient (A) and in thalami in his sibling (B).

1. Finsterer J. Leigh and Leigh-like syndrome in children and adults. *Pediatr Neurol* 2008;39:223–235.
2. Rahman S, Blok RB, Dahl HH, et al. Leigh syndrome: clinical features and biochemical and DNA abnormalities. *Ann Neurol* 1996;39:343–351.
3. Hattori Y, Goto Y, Sakuta R, Nonaka I, Mizuno Y, Horai S. Point mutations in mitochondrial tRNA genes: sequence analysis of chronic progressive external ophthalmoplegia (CPEO). *J Neurol Sci* 1994;125:50–55.
4. Petersen MB, McKusick VA, Smith M. *Online Mendelian Inheritance in Man (OMIM)*. Baltimore: Johns Hopkins University; 2001.
5. Anderson S, Bankier AT, Barrell BG, et al. Sequence and organization of the human mitochondrial genome. *Nature* 1981;290:457–465.
6. Bibb MJ, Van Etten RA, Wright CT, Walberg MW, Clayton DA. Sequence and gene organization of mouse mitochondrial DNA. *Cell* 1981;26:167–180.
7. Kondo R, Horai S, Satta Y, Takahata N. Evolution of hominoid mitochondrial DNA with special reference to the silent substitution rate over the genome. *J Mol Evol* 1993;36:517–531.

Neurology® Genetics

Novel mutation in mitochondrial DNA in 2 siblings with Leigh syndrome

Aravindhan Veerapandiyan, Amit Chaudhari, Christin M. Traba, et al.

Neurol Genet 2016;2;

DOI 10.1212/NXG.0000000000000099

This information is current as of August 16, 2016

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2016 American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/2/5/e99.full.html
References	This article cites 6 articles, 0 of which you can access for free at: http://ng.neurology.org/content/2/5/e99.full.html##ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://ng.neurology.org/content/2/5/e99.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://ng.neurology.org/cgi/collection/all_genetics Cerebellum http://ng.neurology.org/cgi/collection/cerebellum Developmental disorders http://ng.neurology.org/cgi/collection/developmental_disorders Mitochondrial disorders http://ng.neurology.org/cgi/collection/mitochondrial_disorders MRI http://ng.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2016 American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.