

Wei Wang, MD
Yanhong Wu, PhD
Chen Wang, PhD
Jinsong Jiao, MD
Christopher J. Klein, MD

Neurol Genet
2016;2:e117; doi: 10.1212/
NXG.0000000000000117

COPY NUMBER ANALYSIS REVEALS A NOVEL MULTIEXON DELETION OF THE COLQ GENE IN CONGENITAL MYASTHENIA

OPEN

Congenital myasthenic syndrome (CMS) is genetically and clinically heterogeneous.¹ Despite a considerable number of causal genes discovered, many patients are left without a specific diagnosis after genetic testing. The presumption is that novel genes yet to be discovered will account for the majority of such patients. However, it is also possible that we are neglecting a type of genetic variation: copy number changes (>50 bp) as causal for some of these patients. Next-generation sequencing (NGS) can simultaneously screen all known causal genes² and is increasingly being validated to have a potential to identify copy number changes.³ We present a CMS case who did not receive a genetic diagnosis from previous Sanger sequencing, but through a novel copy number analysis algorithm integrated into our targeted NGS panel, we discovered a novel copy number mutation in the *COLQ* gene and made a genetic diagnosis. This discovery expands the genotype-phenotype correlation of CMS, leads to improved genetic counsel, and allows for specific pharmacologic treatment.¹

Case report. The patient is a 16-year-old girl of Han Chinese descent. She had progressive muscle weakness in a limb-girdle pattern and neck weakness since age 6. The muscle weakness appeared more severe after prolonged activity and would partly resolve after minutes of rest. She was seen at the neurology clinic of China-Japan Friendship Hospital in July 2013 and was noted to have a waddling gait and could not walk more than ~600 feet. No ptosis or double vision was present. She had normal sensory testing and was without pain or other sensory symptoms. She had weakness of neck muscles, proximal upper limbs 4/5 (Medical Research Council), and proximal lower limbs 3/5, with normal ankle, hand strengths, and deep tendon reflexes. Her parents and a younger brother did not have any symptoms.


Laboratory testing was unremarkable for creatine kinase, serum lactic acid, anti-acetylcholine receptor, anti-acetylcholinesterase, and anti-Musk antibody. On her EMG, 2 Hz stimulation demonstrated a >10%

decrement without facilitation. Biceps muscle biopsy was without vacuoles, dystrophic changes, or tubular aggregates. Sanger sequencing for *GFPT1*, *DOK-7*, and *COLQ* showed only a heterozygous variant (IVS16+3A→G) in the *COLQ* gene, a previously reported mutation but only deleterious in homozygous state.⁴ This variant was inherited from her father. No genetic diagnosis could be made at that time (figure).

We recently developed a targeted NGS panel including 21 known CMS genes and applied it to this patient. Because a copy number variation (CNV) analysis algorithm (PatternCNV) is incorporated in the bioinformatics evaluation, this panel has a capability of detecting nucleotide changes, small insertion/deletions, and CNVs.³ The result from this panel confirmed the IVS16+3A→G variant and identified a heterozygous copy number deletion encompassing exons 14 and 15 of the *COLQ* gene (~1 kb) (figure, A–C). This copy number change was subsequently confirmed using TaqMan Copy Number Assay (figure, D) and also found in her mother. TaqMan Copy Number Assay was performed using a probe targeting exon 15 of *COLQ* (Hs01393787_cn; Applied Biosystems, Foster City, CA) and housekeeping gene *RNASP* (Applied Biosystems) as copy number control with delta-delta Ct method for copy number calculation. The discovery of this novel copy number mutation of exons 14 and 15 led to the genetic diagnosis of this patient: compound heterozygous *COLQ* mutation, which allowed for more effective genetic counseling in her family. We are initiating drug therapies which have been shown beneficial for patients with *COLQ* mutations.¹

Discussion. Traditional genetic testing often misses large genomic deletions encompassing multiexon (a type of CNV) unless a separate microarray assay is performed. However, comparative genome hybridization arrays would have missed this *COLQ* exons 14 and 15 deletion as the resolution is often inadequate for less than 1 kb deletions. This report underscores the importance of including a validated copy number evaluation in targeted NGS testing in CMS. To date, mutation discovery has been largely focusing on nucleotide changes or small deletion/insertions, mainly due to technology limitations. Current studies estimate that ~10% of the human genome has copy number

Figure Novel compound heterozygous mutation in the *COLQ* gene causes congenital myasthenic syndrome



(A) Pedigree of the proband (II-1). Point mutation (IVS16+3A→G) in the *COLQ* gene was inherited from the father (I-1), and multiexon deletion from the mother (I-2). (B) Gene list of 21 known congenital myasthenic syndrome (CMS) genes included in the targeted CMS panel. (C) Genome view of PatternCNV analysis shows decreased copy number variation (CNV) log₂ ratio for the *COLQ* gene in chromosome 3. (D) Exon-level CNV summary table shows the start and end position of deletion (984 base pairs) in the *COLQ* gene, which indicates 1 copy deletion of exons 14 and 15. SNR.db: signal noise ratio expressed in decibels; CNV.ratio: copy number ratio converted from CNV.log₂ratio. CNV.ratio of 1 indicates no copy number change. (E) TaqMan Copy Number Assay results confirm exon deletions in the *COLQ* gene in the proband, which is also found in her mother (data not shown). Longer PCR cycle number (X-axis) denotes 1 less copy after normalization using *RNASP*.

changes,⁵ a magnitude higher than single nucleotide variants, underlining the importance of understanding copy number changes as we move forward in this molecular diagnostic era. A *COLQ* multiexon deletion (exon 2–3) was first reported once back in 1998.⁶ This is the second multiexon deletion of *COLQ* discovered in almost 20 years. With qPCR method, multiexon deletions of *RAPSN* were recently reported in 15% of previously undiagnosed patients with CMS,⁷ emphasizing the pathogenic role of copy number changes. Validating and providing copy number evaluation in targeted NGS will considerably improve the efficacy of genetic testing and is predicted to reduce the overall cost for CMS and possibly for other genetic neuromuscular disorders.

From the Department of Neurology (W.W., J.J.), China-Japan Friendship Hospital, Beijing, China; Department of Neurology (W.W., C.J.K.), Department of Laboratory Medicine and Pathology (Y.W.), Department of Health Sciences Research (C.W.), and Department of Medical Genetics (C.J.K.), Mayo Clinic, Rochester, MN.

Author contributions: Dr. Wei Wang: study concept and design, acquisition of data, analysis and interpretation, drafting the manuscript, and critical revision of the manuscript. Dr. Yanhong Wu: acquisition of data, analysis and interpretation, and critical revision of the manuscript. Dr. Chen Wang: analysis and interpretation of data and critical revision of the manuscript. Dr. Jinsong Jiao: study concept and design, acquisition of data, analysis and interpretation, and critical revision of the manuscript. Dr. Christopher J. Klein: study concept and design; acquisition of data, analysis and interpretation, drafting the manuscript, critical revision of the manuscript, study supervision, and laboratory funding and support.

Study funding: No targeted funding reported.

Disclosure: Dr. W. Wang, Dr. Wu, Dr. C. Wang, and Dr. Jiao report no disclosures. Dr. Klein has served on the editorial board of the Journal of Peripheral Nerve Society and has received research support from a Mayo Clinic research grant. Go to Neurology.org/ng for full disclosure forms. The Article Processing Charge was paid by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Received July 26, 2016. Accepted in final form September 22, 2016.

Correspondence to Dr. Klein: klein.christopher@mayo.edu or Dr. Jiao: jiaojinsong@alinyun.com

1. Engel AG, Shen XM, Selcen D, Sine SM. Congenital myasthenic syndromes: pathogenesis, diagnosis, and treatment. *Lancet Neurol* 2015;14:461.
2. Abicht A, Dusl M, Gallenmuller C, et al. Congenital myasthenic syndromes: achievements and limitations of phenotype-guided gene-after-gene sequencing in diagnostic practice: a study of 680 patients. *Hum Mutat* 2012;33:1474–1484.
3. Wang W, Wang C, Dawson DB, et al. Target-enrichment sequencing and copy number evaluation in inherited polyneuropathy. *Neurology* 2016;86:1762–1771.
4. Ohno K, Brengman JM, Felice KJ, Cornblath DR, Engel AG. Congenital end-plate acetylcholinesterase deficiency caused by a nonsense mutation and an A→G splice-donor-site mutation at position +3 of the collagenlike-tail-subunit gene (COLQ): how does G at position +3 result in aberrant splicing? *Am J Hum Genet* 1999;65:635–644.
5. Zarrei M, MacDonald JR, Merico D, Scherer SW. A copy number variation map of the human genome. *Nat Rev Genet* 2015;16:172–183.
6. Ohno K, Brengman J, Tsujino A, Engel AG. Human end-plate acetylcholinesterase deficiency caused by mutations in the collagen-like tail subunit (ColQ) of the asymmetric enzyme. *Proc Natl Acad Sci USA* 1998;95:9654–9659.
7. Gaudon K, Penisson-Besnier I, Chabrol B, et al. Multiexon deletions account for 15% of congenital myasthenic syndromes with RAPSN mutations after negative DNA sequencing. *J Med Genet* 2010;47:795–796.

Neurology[®] Genetics

Copy number analysis reveals a novel multiexon deletion of the *COLQ* gene in congenital myasthenia

Wei Wang, Yanhong Wu, Chen Wang, et al.

Neurol Genet 2016;2;

DOI 10.1212/NXG.0000000000000117

This information is current as of October 31, 2016

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/2/6/e117.full.html
References	This article cites 7 articles, 2 of which you can access for free at: http://ng.neurology.org/content/2/6/e117.full.html##ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://ng.neurology.org/content/2/6/e117.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://ng.neurology.org/cgi/collection/all_genetics All Neuromuscular Disease http://ng.neurology.org/cgi/collection/all_neuromuscular_disease Myasthenia http://ng.neurology.org/cgi/collection/myasthenia
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2016 American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.

