Rubén Fernández-
Santiago, PhD*
Alex Iranzo, MD*
Carles Gaig, MD
Mónica Serradell, BSc
Manel Fernández
Pau Pastor, MD
Eduardo Tolosa, MD
Joan Santamaría, MD
Mario Ezquerro, PhD

Neurol Genet
2017;3:e131; doi: 10.1212/
NXG.0000000000000131

MAPT ASSOCIATION WITH REM SLEEP BEHAVIOR DISORDER

OPEN

Idiopathic REM sleep behavior disorder (IRBD) is a REM sleep parasomnia comprising unpleasant dreams, dream-enacting behaviors, and loss of muscle atonia during REM sleep. Longitudinal studies have demonstrated that because most patients with IRBD develop with time Parkinson disease (PD) and other synucleinopathies including dementia with Lewy bodies (DLB) or multiple system atrophy (MSA), IRBD represents a prodromal stage of these diseases.¹ In PD, 5%–10% of cases are caused by nonfrequent mendelian mutations segregating with disease in pedigrees, but the vast majority of cases are sporadic. Unbiased genome-wide association studies in sporadic PD (sPD) have shown that single nucleotide polymorphisms (SNPs) in the α -synuclein (*SNCA*) and microtubule-associated protein tau (*MAPT*) genes modulate disease susceptibility.² Given that IRBD often antedates sPD and that, akin to PD, familial clustering of IRBD is rare, we hypothesized that genetically shared variation at *SNCA* and *MAPT* may influence disease susceptibility to both conditions. Accordingly, we genotyped PD-associated genetic variants in *SNCA* and *MAPT* in a cohort of Spanish patients with IRBD.

Methods. The SNPs rs356219 (A/G) in *SNCA* and rs1800547 (H1/H2) in *MAPT* have been consistently associated with sPD in the Spanish population.^{3,4} Here, we genotyped these markers in a Spanish Caucasian sample of 121 patients with IRBD and 175 healthy controls. This population was studied in a previous report.⁵ The IRBD group was recruited at the Multidisciplinary Sleep Unit of the Hospital Clínic de Barcelona, had a polysomnography-confirmed diagnosis, and consisted of 21/100 women/men (17.4/82.6%), with age at IRBD onset of 68.3 ± 6.2 years and age at sample collection of 71.2 ± 6.4 years. Patients fulfilled the diagnosis of IRBD and were free of parkinsonism, mild cognitive impairment, and dementia at the time of sample collection. After 6.8 ± 4.1 years of follow-up from IRBD diagnosis, 38 patients (31.4%) were clinically diagnosed with a synucleinopathy (DLB in 19

patients, PD in 18, and MSA in 1). Sex-, age-, and demographic-matched controls were recruited at the Multidisciplinary Sleep Unit of the Hospital Clínic de Barcelona (n = 45) and the Clínica Universitaria de Navarra (n = 129), and consisted of 21/154 women/men (13.6/86.4%) with age at sample collection of 71.7 ± 6.4 years. DNA was extracted from the peripheral blood using standard procedures. Genotyping was performed using the predesigned TaqMan assays C-1020193-10 (*SNCA* rs356219) and C-7563692-10 (*MAPT* rs1800547) in a StepOnePlus Real-time PCR System (Applied Biosystems, Foster City, CA). Allelic and genotypic frequencies were compared using SNPstats software (bioinfo.iconcologia.net/SNPstats).

Standard protocol approvals, registrations, and patient consents. Samples were collected after patient's signed consent, and the local ethics committee approved the study.

Results. We found no allelic or genotypic distribution differences of the *SNCA* marker rs356219 in patients with IRBD and in controls. However, we found an association of the *MAPT* polymorphism rs1800547 with IRBD. More specifically, we found that the frequency of the *MAPT* H2/H2 genotype which is protective for PD^{3,4} was underrepresented in our patients with IRBD. This association held statistically significant after adjusting for sex and age as covariates (odds ratio = 0.16; 95% confidence interval = 0.04–0.69; $p = 0.002$; table). Moreover, we observed that *MAPT* genotypes in patients with IRBD, but not in controls, were not in the Hardy-Weinberg equilibrium ($p = 0.049$), a finding which is consistent with previous observations for disease-linked loci in disease-affected individuals.⁶

Discussion. We found that the *MAPT* marker rs1800547 is associated with IRBD. Consistently, we detected that the H2 variant which is protective for PD was underrepresented in our patients with IRBD. On the contrary, we did not find an association of the *SNCA* polymorphism rs356219 with IRBD. Altogether, these results are in agreement with another IRBD study reporting association for *MAPT* but not for *SNCA* polymorphisms.⁷ Yet, markers in both studies were different. These data raise the question whether IRBD shares with PD the same genetic

Table Distribution of genotypic frequencies of SNCA and MAPT polymorphisms in patients with IRBD (n = 121) and sex- and age-matched healthy controls (n = 175)

Marker	Frequency in IRBD	Frequency in controls	OR (95% CI)	Adjusted p value
MAPT rs1800547				
H1/H1	0.537 (n = 65)	0.474 (n = 83)	0.16 (0.04–0.69)	0.002
H1/H2	0.438 (n = 53)	0.423 (n = 74)		
H2/H2	0.025 (n = 3)	0.103 (n = 18)		
SNCA rs356219				
A/A	0.562 (n = 68)	0.497 (n = 87)	0.45 (0.31–1.69)	0.45
A/G	0.364 (n = 44)	0.400 (n = 70)		
G/G	0.074 (n = 9)	0.103 (n = 18)		

Abbreviations: CI = confidence interval; IRBD = idiopathic REM sleep behavior disorder; OR = odds ratio.

p Values were calculated using the Student t test, adjusting for sex and age as covariates, and comparing the MAPT PD-protective genotype H2/H2 vs H1/H1 + H1/H2 and the SNCA PD-risk SNCA genotype G/G vs G/A + A/A.

risk factors or whether, alternatively, IRBD represents a specific endophenotype of PD with only certain genetic risk factors shared in common by both conditions. The later might be a plausible explanation because, although IRBD is considered a prodrome of sPD, not all patients with sPD develop RBD prior to parkinsonism. Supporting this view, we have recently reported that pathogenic mutations in the leucine-rich repeat kinase 2 gene (*LRRK2*), which are the most frequent genetic cause of PD, are absent in our IRBD cohort.⁵ One limitation of our study is the reduced number of patients with IRBD. Thus, future studies in larger cohorts are warranted to validate the association of *MAPT* with IRBD and also to further elucidate whether or not *SNCA* polymorphisms play a role in disease. We found that one polymorphism in *MAPT* which is associated with PD also modulates the propensity to IRBD. These results point toward an at least in part overlapping genetic susceptibility to both conditions in our Spanish population.

*These authors contributed equally to this work.

From the Laboratory of Neurodegenerative Disorders (R.F.-S., M.F., E.T., M.E.), Multidisciplinary Sleep Unit (A.I., C.G., M.S., J.S.), Movement Disorders Unit (E.T.), Department of Neurology, Hospital Clínic de Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona; Centro de Investigación Biomédica en Red de Enfermedades Neurodegenerativas (CIBERNED) (R.F.-S., A.I., C.G., M.S., M.F., E.T., J.S., M.E.), Madrid; and Movement Disorders Unit (P.P.), Department of Neurology, University Hospital Mutua de Terrassa, Barcelona, Spain.

Author contributions: R.F.-S., A.I., and M.E. codirected the study and wrote the first draft of the manuscript. A.I., C.G., P.P., M.S., E.T., and J.S. diagnosed and followed up the patients, and recruited the controls. R.F.-S., M.F., P.P., and M.E. performed the genotyping. All authors interpreted the data and revised the manuscript.

Acknowledgment: The authors thank the patients and their relatives who participated in the study. This study was conducted at the Centre de Recerca Biomèdica Cellex, Barcelona, Spain. The authors also

thank the CERCA from the Generalitat de Catalunya and the European FEDER Programmes to IDIBAPS.

Study funding: This work was supported by funds from the Centro de Investigación Biomédica en Red de Enfermedades Neurodegenerativas (CIBERNED) to the Movement Disorders Unit of the Neurology Service from the Hospital Clínic de Barcelona (grant PRI-16-207 to R.F.-S., A.I., C.G., M.S., M.F., E.T., J.S., and M.E.), and from the Spanish Ministry of Economy and Competitiveness (MINECO) (grant SAF2013-47939-R to P.P.). R.F.-S. was supported by a Marie Skłodowska-Curie contract of the European Commission (EC) and Juan-de-la-Cierva contract of the Spanish Ministry of Economy and Competitiveness (MINECO), and M.E. by a Miguel Servet contract of the Instituto de Salud Carlos III (ISCIII).

Disclosure: Dr. Fernández-Santiago, Dr. Iranzo, Dr. Gaig, Ms. Serradell, Mr. Fernández, and Dr. Pastor report no disclosures. Dr. Tolosa has received speaker honoraria from UCB, Boehringer Ingelheim, Novartis, Abbot, Medtronic, Solvay, GSK, and Teva; has served on the editorial board of Neurological Sciences; holds a patent for Method for the subclassification of patients suffering from Parkinson disease; and has been a consultant for the Michael J. Fox Foundation for Parkinson's Disease, Novartis, UCB, Abbot, and Teva. Dr. Santamaría has received research support from Fundació La Marató TV3. Dr. Ezquerra reports no disclosures. Go to Neurology.org for full disclosure forms. The Article Processing Charge was paid by CIBERNED.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Received September 16, 2016. Accepted in final form October 28, 2016.

Correspondence to Dr. Fernández-Santiago: ruben.fernandez.santiago@gmail.com or Dr. Ezquerra: ezquerra@clinic.ub.es

1. Iranzo A, Tolosa E, Gelpi E, et al. Neurodegenerative disease status and post-mortem pathology in idiopathic rapid-eye-movement sleep behaviour disorder: an observational cohort study. *Lancet Neurol* 2013;12:443–453.
2. Simon-Sanchez J, Schulte C, Bras JM, et al. Genome-wide association study reveals genetic risk underlying Parkinson's disease. *Nat Genet* 2009;41:1308–1312.
3. Pastor P, Ezquerra M, Munoz E, et al. Significant association between the tau gene A0/A0 genotype and Parkinson's disease. *Ann Neurol* 2000;47:242–245.

4. Botta-Orfila T, Ezquerro M, Rios J, et al. Lack of interaction of SNCA and MAPT genotypes in Parkinson's disease. *Eur J Neurol* 2011;18:e32.
5. Fernandez-Santiago R, Iranzo A, Gaig C, et al. Absence of LRRK2 mutations in a cohort of patients with idiopathic REM sleep behavior disorder. *Neurology* 2016;86:1072–1073.
6. Lee WC. Searching for disease-susceptibility loci by testing for Hardy-Weinberg disequilibrium in a gene bank of affected individuals. *Am J Epidemiol* 2003;158:397–400.
7. Gan-Or Z, Girard SL, Noreau A, et al. Parkinson's disease genetic loci in rapid eye movement sleep behavior disorder. *J Mol Neurosci* 2015;56:617–622.

Neurology[®] Genetics

***MAPT* association with REM sleep behavior disorder**
Rubén Fernández-Santiago, Alex Iranzo, Carles Gaig, et al.
Neurol Genet 2017;3;
DOI 10.1212/NXG.0000000000000131

This information is current as of January 13, 2017

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/3/1/e131.full.html
References	This article cites 7 articles, 0 of which you can access for free at: http://ng.neurology.org/content/3/1/e131.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Dementia with Lewy bodies http://ng.neurology.org/cgi/collection/dementia_with_lewy_bodies Parkinson's disease/Parkinsonism http://ng.neurology.org/cgi/collection/parkinsons_disease_parkinsonism
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2017 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.