Gaël Nicolas, MD, PhD
Florent Marguet, MD
Annie Laquerrière, MD,
PhD
João Ricardo Mendes de
Oliveira, MD, PhD
Didier Hannequin, MD,
PhD

Neurol Genet
2017;3:e134; doi: 10.1212/
NXG.0000000000000134

MICROANGIOPATHY IN PRIMARY FAMILIAL BRAIN CALCIFICATION: EVIDENCE FROM SKIN BIOPSIES

OPEN

Autosomal dominant primary familial brain calcification (PFBC) is a rare cerebral microvascular calcifying disorder defined by the presence of calcifications affecting at least the basal ganglia with no secondary cause. It is associated with diverse symptoms including movement disorders, psychiatric disturbances, and cognitive impairment.¹ PFBC is caused by loss-of-function mutations in 2 groups of genes: (1) *PDGFB*, which encodes the platelet-derived growth factor B² and *PDGFRB*, which encodes its main receptor platelet-derived growth factor receptor- β (PDGFR- β)³ and (2) *SLC20A2* and *XPR1* encoding inorganic phosphate transporters.^{4,5} Mice carrying *Pdgfb* hypomorphic alleles exhibit lower pericyte coverage in cerebral microvessels, blood-brain barrier (BBB) impairment, and cerebral microvascular calcifications.² Recently, a novel *PDGFB* mutation was reported in an Italian family with PFBC and white-matter hyperintensities (WMH).⁶ Although brain calcification is a mandatory criterion for diagnosing PFBC, WMH were also reported as a major neuroimaging feature in the first described families with a *PDGFRB* or a *PDGFB* mutation^{2,3} (table e-1 at Neurology.org/ng). To date, the precise nature of WMH remains unknown but may be regarded as resulting from microangiopathy. This led to the hypothesis that in mice, alterations of the microvessels leading to BBB impairment may be a causal mechanism between microangiopathy and vascular calcifications.² Transmission electron microscopy analysis of a skin biopsy from a patient belonging to the above-mentioned *PDGFB* family revealed thickened and fragmented areas in the basal lamina, consistent with microangiopathy.⁶ We report here the results of skin biopsies performed in 2 patients carrying a *PDGFRB* and an *XPR1* mutation, respectively.

Methods. This study was approved by our institution's ethics committee. After having obtained written informed consent from the patients, punch biopsy was performed. Ultrastructural studies were conducted according to standardized protocols. Briefly, tissue

samples were fixed in a 2% glutaraldehyde fixative solution, postfixed with osmium tetroxide, and embedded in resin epoxy. Semithin sections were stained with toluidine blue. Ultrathin sections were contrasted with uranyl acetate and lead citrate and examined under a Philips CM10 electron microscope.

Results. In the proband of the family with the p.Leu658Pro *PDGFRB* mutation,³ biopsy analysis revealed no lesions of endothelial cells, whereas the basal lamina was thickened, and with microcalcifications within and around the pericytes and in the basal lamina (figure, A). These microcalcifications were sometimes included in double- or single-layered membranes in the vicinity of the basal lamina (figure, B). In another patient with a recently described p.Ser136Asn *XPR1* mutation,⁵ microcalcifications were also observed within pericytes of the capillaries (figure, C) but remained located in the cytoplasm of the pericytes, laying under the plasma membrane (figure, D).

Discussion. We herein report hypodermal microvessel calcifications in skin biopsies from patients with PFBC. To our knowledge, only one skin biopsy analysis was previously reported in a patient with PFBC, who carried a *PDGFB* mutation.⁶ As for the latter patient, the basal lamina appeared thickened in our patients, particularly in the *PDGFRB* mutation carrier, although no fragmentation was observed, indicating that *PDGFRB* and *XPR1* mutation carriers also exhibit microangiopathy. WMH on fluid-attenuated inversion recovery or T2-weighted MRI have also been observed in *SLC20A2* and *XPR1* mutation carriers (table e-1). These lesions are therefore not specific to *PDGFB* or *PDGFRB* mutation carriers in which BBB alteration was thought to be a prominent disease mechanism. Furthermore, similar thickening of the microvessel basal lamina on skin biopsies has also been observed in other leukoencephalopathies such as *COL4A1*-related disorders,⁷ where different patterns of calcifications might be encountered in some cases. A summary of the literature review of vascular, clinical, imaging, and microvessel examination of skin biopsies found in PFBC and 2 other leukoencephalopathies (cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy and *COL4A1*-related disorders) is provided in table e-1.

Supplemental data at
Neurology.org/ng

Figure Ultrastructural hallmarks of small capillary lesions in the 2 patients

p.Leu658Pro *PDGFRB* mutation carrier (A, B): presence of microcalcifications in a pericyte (A, white arrow) and in the basal lamina (A, black arrow) (OM $\times 5,200$), appearing to be membrane bound at a higher magnification (OM $\times 28,500$) (B). p.Ser136Asn *XPR1* mutation carrier (C, D): small calcifications in the pericytes (C, white arrows) (OM $\times 2,650$), sometimes located under the pericyte plasma membrane (OM $\times 28,500$) (D). L = lumen; E = endothelial cell; B = basal lamina; P = pericyte; OM = original magnification.

Microangiopathy likely results from different mechanisms in the 2 groups of patients with PFBC and in other cerebral microangiopathies.

Whether BBB alteration is the cause of calcification in *PDGFB* or *PDGFRB* mutation carriers or not is currently debated. The original group who linked BBB deficiency in *PDGFB*-deficient mice and in humans with mutations in the same gene has recently shown that calcification-prone regions in *Pdgfr^{ret/ret}* mice had a more intact BBB and higher pericyte coverage compared with calcification-nonprone brain regions.⁸ Additional studies in *Slc20a2* knockout mice suggest that brain calcifications are found even with normal BBB structure and function, through a 2-hit mechanism whereby increased CSF inorganic phosphate leads to calcification in arteriolar smooth muscle cells due to an enhanced vulnerability caused by *Slc20a2* deficiency.⁹ Although the existence of microangiopathy itself in the context of *PDGFB* or *PDGFRB* haploinsufficiency is not challenged by the recent mouse model report,⁸ a putative direct causal link between microangiopathy and brain calcification is highly questioned. Additional

studies on mice models might also evaluate the existence of microangiopathy outside the brain and if these models reproduce properly the phenotype variability found in patients.

Microvascular changes on skin biopsy are not specific to *PDGFB* mutation carriers. Systematic examination of skin biopsies of other patients with PFBC or differential diagnoses is warranted to replicate and explore these observations in depth. The significance of microangiopathy in both groups of patients and the mechanisms leading to microvascular changes in *XPR1* or *SLC20A2* mutation carriers remain to be determined, but further encourage to search for the potential pathways connecting the PDGFB/PDGFR- β response to the inorganic phosphate transporters *SLC20A2* and *XPR1*.

From the Normandie Univ (G.N., F.L., A.L., D.H.), UNIROUEN, Inserm U1245; Department of Genetics (G.N., D.H.), Department of Pathology (F.M., A.L.), Department of Neurology (D.H.), and CNR-MAJ (G.N., D.H.), Rouen University Hospital, Normandy Center for Genomic and Personalized Medicine, Rouen, France; and Keizo Asami Laboratory (LIKA) (J.R.M.d.O.), and Neuropsychiatry Department (J.R.M.d.O.), Universidade Federal de Pernambuco, Recife, Brazil.

Author contributions: Collection and interpretation of data: Gaël Nicolas, Florent Marguet, Annie Laquerrière, and Didier Hannequin. Manuscript draft: Gaël Nicolas, João Ricardo Mendes de Oliveira, Florent Marguet, and Annie Laquerrière. Critical revision: Gaël Nicolas, Florent Marguet, Annie Laquerrière, Didier Hannequin, and João Ricardo Mendes de Oliveira. Study design and supervision: Gaël Nicolas.

Study funding: No targeted funding reported.

Disclosure: Dr. Nicolas has received research support from Rouen University Hospital and Région Haute Normandie. Dr. Marguet and Dr. Laquerrière report no disclosures. Dr. Mendes de Oliveira has received research support from the Federal Council for Research and Technology and Brazil (CNPq) Foundation for research support in Pernambuco, Brazil (FACEPE). Dr. Hannequin has received research support from the French Ministry of Health. Go to Neurology.org/ng for full disclosure forms. The Article Processing Charge was paid by Inserm.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Received November 19, 2016. Accepted in final form December 29, 2016.

Correspondence to Dr. Nicolas: gaelnicolas@hotmail.com

1. Nicolas G, Charbonnier C, de Lemos RR, et al. Brain calcification process and phenotypes according to age and sex: lessons from *SLC20A2*, *PDGFB*, and *PDGFRB* mutation carriers. *Am J Med Genet B Neuropsychiatr Genet* 2015;168:586–594.
2. Keller A, Westenberger A, Sobrido MJ, et al. Mutations in the gene encoding PDGF-B cause brain calcifications in humans and mice. *Nat Genet* 2013;45:1077–1082.
3. Nicolas G, Pottier C, Maltete D, et al. Mutation of the *PDGFRB* gene as a cause of idiopathic basal ganglia calcification. *Neurology* 2013;80:181–187.

4. Wang C, Li Y, Shi L, et al. Mutations in SLC20A2 link familial idiopathic basal ganglia calcification with phosphate homeostasis. *Nat Genet* 2012;44:254–256.
5. Legati A, Giovannini D, Nicolas G, et al. Mutations in XPR1 cause primary familial brain calcification associated with altered phosphate export. *Nat Genet* 2015;47:579–581.
6. Biancheri R, Severino M, Robbiano A, et al. White matter involvement in a family with a novel PDGFB mutation. *Neurol Genet* 2016;2:e77.
7. Plaisier E, Gribouval O, Alamowitch S, et al. COL4A1 mutations and hereditary angiopathy, nephropathy, aneurysms, and muscle cramps. *N Engl J Med* 2007; 357:2687–2695.
8. Vanlandewijck M, Lebouvier T, Andaloussi Mae M, et al. Functional characterization of germline mutations in PDGFB and PDGFRB in primary familial brain calcification. *PLoS One* 2015;10:e0143407.
9. Wallingford MC, Chia J, Leaf EM, et al. SLC20A2 deficiency in mice leads to elevated phosphate levels in cerebrospinal fluid and glymphatic pathway-associated arteriolar calcification, and recapitulates human idiopathic basal ganglia calcification. *Brain Pathol* 2017;27:64–76.

Neurology[®] Genetics

Microangiopathy in primary familial brain calcification: Evidence from skin biopsies

Gaël Nicolas, Florent Marguet, Annie Laquerrière, et al.

Neurol Genet 2017;3;

DOI 10.1212/NXG.0000000000000134

This information is current as of February 8, 2017

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/3/2/e134.full.html
Supplementary Material	Supplementary material can be found at: http://ng.neurology.org/content/suppl/2017/02/08/3.2.e134.DC1
References	This article cites 9 articles, 1 of which you can access for free at: http://ng.neurology.org/content/3/2/e134.full.html##ref-list-1
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2017 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.