Mateja Smogavec, MD*
Jana Zschüntzsch, MD*
Wolfram Kress, MD
Julia Mohr, PhD
Peter Hellen, MD
Barbara Zoll, MD
Silke Pauli, MD†
Jens Schmidt,
MD, FEAN†

Neurol Genet
2017;3:e167; doi: 10.1212/
NXG.0000000000000167

NOVEL FUKUTIN MUTATIONS IN LIMB-GIRDLE MUSCULAR DYSTROPHY TYPE 2M WITH CHILDHOOD ONSET

OPEN

Limb-girdle muscular dystrophies (LGMDs) are a heterogeneous group of childhood- or adult-onset inherited neuromuscular disorders, which are characterized by weakness and wasting of proximal limb and axial muscles. LGMDs are divided into 8 autosomal-dominant (LGMD1)¹ and 25 autosomal-recessive (LGMD2) forms (omim.org/phenotypicSeries/PS253600). One autosomal-recessive form is the LGMD2M (new nomenclature: Muscular Dystrophy-Dystroglycanopathy, type C, 4, MDDGC4, MIM #611588), which is caused by homozygous or compound heterozygous mutations in the fukutin gene (*FKTN*; MIM #607440) on chromosome 9q31.2. The LGMD2M, as the clinical mild end of fukutin-related muscle dystrophy without mental retardation, is rare worldwide and described mostly in individuals of non-Japanese descent (table e-1 at Neurology.org/ng).²⁻⁵

Here, we report the second German case with an LGMD2M phenotype and describe 2 previously unreported mutations in the *FKTN* gene in a German female patient.

Clinical case presentation. A 32-year-old female patient was admitted to our Neuromuscular Center. Written consent was obtained for the use of all clinical and diagnostic data. The patient was born to healthy nonconsanguineous parents. At the age of 7 years, cramping muscle pain, especially in the thighs, started and was accompanied by creatine kinase (CK) elevation. A muscle biopsy record described a chronic myopathic pattern and marked inflammation. The patient was treated with a corticosteroid, which profoundly reduced the CK level. In the following years, the patient developed a slowly progressive muscle weakness of both legs. At the age of 16 years, problems in climbing stairs and lifting heavy objects were noted. At the time of admission, the patient started to use a wheelchair for outdoor activities.

Neurologic examination revealed normal intellectual status, cranial nerve function, and sensorium. Inspection showed scapular winging, calf muscle enlargement, toe walking, and lumbar hyperlordosis

(figure, A and D). We observed muscular atrophy with accentuation of the proximal lower extremities and reduced muscle strength. Gower and Trendelenburg signs were positive. Deep tendon reflexes were diminished. The patient also suffered from mild dysphagia and respiratory distress. EMG showed myopathic changes. Muscular MRI revealed a pronounced lipodystrophy of proximal upper and lower limb and limb-girdle muscles (figure, B, C, E, G, and H).

A multigene panel for LGMDs by next-generation sequencing (NGS) on genomic DNA isolated from a blood sample of our patient was performed. Coding and flanking sequences of all relevant genes were enriched using Agilent SureSelect technology. Massively parallel sequencing was performed using the HiSeq 2500 Sequencing platform from Illumina. Sanger sequencing was used for badly covered regions and validation of potentially pathogenic mutations.

In our patient, 2 novel missense mutations c.895A>C; p.Ser299Arg and c.1325A>G; p.Asn442Ser of the *FKTN* gene (NM_001079802.1) were found by NGS analysis and confirmed by Sanger sequencing. Both mutations disturb highly conserved positions in the fukutin protein. Several in silico prediction programs (MutationTaster, PolyPhen-2, SIFT) indicated that these mutations are most likely damaging. In Exome Aggregation Consortium (ExAC) and in Exome Server Browser, the variant c.895A>C was detected in heterozygous state only in one person, whereas the variant c.1325A>G had not been observed before.

With only one mutation c.1325A>G present in the father, a compound heterozygosity of the 2 mutations in our patient is strongly implied. Based on all information, we predict these 2 mutations as causative for the patients' disease.

Discussion. Mutations in *FKTN* cause a wide clinical spectrum of myopathies: from severe congenital forms of muscular dystrophy with additional extramuscular symptoms to a milder form of LGMD2M, which is normally not associated with cognitive impairment.² In the phenotypical spectrum of *FKTN* mutations, the classification from congenital muscular dystrophy (CMD) to LGMD2M was defined according to the onset of weakness.² In practice, patients might display a phenotype well in between CMD and LGMD,^{2,6}

Supplemental data at
Neurology.org/ng

Clinical presentation: atrophy of the proximal muscles in the upper extremities. Note the scapula alata (A) and the characteristic symmetric hypertrophy of gastrocnemius muscles (D). Imaging findings: MRI of the upper extremity demonstrates a moderate atrophy of the rotator cuff and the deltoid muscle (B and C). MRI of the lower extremity shows a severe symmetric atrophy with fatty degeneration of musculature of both thighs (E and F). The less atrophied sartorius and gracilis muscles depict a pronounced edema (G) and a strong gadolinium enhancement representing acute disease activity (H and I). B, C, E, and F: transverse T1-weighted images, G: coronal STIR sequence; H and I: transverse fat-saturated T1-weighted images postcontrast.

pointing to an inherent difficulty of a strict classification. The phenotype of our patient is characterized by childhood onset, hypertrophy of calves, normal intelligence, and initial responsiveness to steroids.

The reduced ability of our patient to walk in her teens is in line with previous reports.⁷ In addition, we can provide clinical data of our patient until the age of 32 years. Our patient initially responded to steroid therapy, which might be explained by the profound inflammation in the muscle biopsy, comparable with the positive effect of steroids in Duchenne muscular dystrophy.

We show a rare case of LGMD2M and provide further insight into heterogeneity of phenotypes caused by mutations in *FKTN* and underline the importance of broad genetic assessment to provide correct diagnosis and to facilitate individual treatment in the future.

*These authors contributed equally to this work as co-first authors.

†These authors contributed equally to this work as co-last authors.

From the Institute of Human Genetics (M.S., B.Z., S.P.), Department of Neurology (J.Z., J.S.), and Department of Neuroradiology (P.H.), University Medical Center Göttingen; Department of Human Genetics (W.K.), University of Würzburg; and CeGaT GmbH und Praxis für Humangenetik (J.M.), Tübingen, Germany.

Author contributions: Mateja Smogavec and Jana Zschüntzsch: case report study and concept and design. Wolfram Kress: analysis and interpretation of data and critical revision of manuscript. Julia Mohr: analysis and interpretation of data. Peter Hellen: interpretation of radiologic imaging. Barbara Zoll: critical revision of the manuscript. Silke Pauli and Jens Schmidt: case report study concept and critical revision of the manuscript.

Study funding: No targeted funding reported.

Disclosure: M. Smogavec reports no disclosures. J. Zschüntzsch has received travel funding/speaker honoraria from Biotest AG and Grifols. W. Kress reports no disclosures. J. Mohr has been an employee of CeGaT GmbH and Praxis für Humangenetik. P. Hellen, B. Zoll, and S. Pauli report no disclosures. J. Schmidt has served on scientific advisory boards for BioMarin and CSL Behring; has received travel funding/speaker honoraria from Bayer, Biogen, Biotest, CSL Behring, Grifols, Novartis, Octapharma, and VitalAire; has served on the editorial boards of BMC Neurology, American Journal of Neuroscience, Edorium Journal of Neurology, World Journal of Neurology, and Journal of Neuromuscular Diseases; has been a consultant for Novartis and CSL Behring; and has received research support from Biogen, CSL Behring, Novartis, Octapharma, Association française contre les myopathies, and Deutsche Gesellschaft für Muskelkranke. Go to Neurology.org/ng for full disclosure forms. The Article Processing Charge was funded by Institute of Human Genetics Göttingen.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Received February 8, 2017. Accepted in final form April 28, 2017.

Correspondence to Dr. Smogavec: mateja.smogavec@med.uni-goettingen.de or Prof. Schmidt: j.schmidt@gmx.org

1. Nigro V, Savarese M. Genetic basis of limb-girdle muscular dystrophies: the 2014 update. *Acta Myol* 2014;33:1–12.
2. Godfrey C, Clement E, Mein R, et al. Refining genotype phenotype correlations in muscular dystrophies with defective glycosylation of dystroglycan. *Brain* 2007;130:2725–2735.

3. Vuillaumier-Barrot S, Quijano-Roy S, Bouchet-Seraphin C, et al. Four Caucasian patients with mutations in the fukutin gene and variable clinical phenotype. *Neuromuscul Disord* 2009;19:182–188.
4. Saredi S, Ruggieri A, Mottarelli E, et al. Fukutin gene mutations in an Italian patient with early onset muscular dystrophy but no central nervous system involvement. *Muscle Nerve* 2009;39:845–848.
5. Puckett RL, Moore SA, Winder TL, et al. Further evidence of Fukutin mutations as a cause of childhood onset limb-girdle muscular dystrophy without mental retardation. *Neuromuscul Disord* 2009;19:352–356.
6. Ceyhan-Birsoy O, Talim B, Swanson LC, et al. Whole exome sequencing reveals DYSE, FKTN, and ISPD mutations in congenital muscular dystrophy without brain or eye involvement. *J Neuromuscul Dis* 2015;2:87–92.
7. Riisager M, Duno M, Hansen FJ, Krag TO, Vissing CR, Vissing J. A new mutation of the fukutin gene causing late-onset limb girdle muscular dystrophy. *Neuromuscul Disord* 2013;23:562–567.

Neurology[®] Genetics

Novel fukutin mutations in limb-girdle muscular dystrophy type 2M with childhood onset

Mateja Smogavec, Jana Zschüntzsch, Wolfram Kress, et al.

Neurol Genet 2017;3;

DOI 10.1212/NXG.0000000000000167

This information is current as of July 10, 2017

Updated Information & Services

including high resolution figures, can be found at:
<http://ng.neurology.org/content/3/4/e167.full.html>

Supplementary Material

Supplementary material can be found at:
<http://ng.neurology.org/content/suppl/2017/07/10/3.4.e167.DC1>

References

This article cites 7 articles, 0 of which you can access for free at:
<http://ng.neurology.org/content/3/4/e167.full.html##ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Muscle disease
http://ng.neurology.org/cgi/collection/muscle_disease

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://ng.neurology.org/misc/about.xhtml#permissions>

Reprints

Information about ordering reprints can be found online:
<http://ng.neurology.org/misc/addr.xhtml#reprintsus>

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2017 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.