Psychometric properties of the Friedreich Ataxia Rating Scale

Christian Rummey, PhD, Louise A. Corben, PhD, Martin B. Delatycki, MD, S.H. Subramony, MD, Khalaf Bushara, MD, Christopher M. Gomez, MD, Joseph Chad Hoyle, MD, Grace Yoon, MD, Bernard Ravina, MD, Katherine D. Mathews, MD, George Wilmot, MD, Theresa Zesiewicz, MD, Susan Perlman, MD, Jennifer M. Farmer, MD, and David R. Lynch, MD, PhD

Correspondence

Dr. Lynch
lynchd@mail.med.upenn.edu

Neurol Genet 2019;5:e371. doi:10.1212/NXG.0000000000000371

Abstract

Objective

To investigate the psychometric properties of the Friedreich Ataxia Rating Scale neurologic examination (FARSn) and its subscores, as well as the influence of the modifications resulting in the now widely used modified FARS (mFARS) examination.

Methods

Based on cross-sectional FARS data from the FA–Clinical Outcome Measures cohort, we conducted correlation-based psychometric analyses to investigate the interplay of items and subscores within the FARSn/mFARS constructs.

Results

The results provide support for both the FARSn and the mFARS constructs, as well as individually for their upper limb and lower limb coordination components. The omission of the peripheral nervous system subscore (D) and 2 items of the bulbar subscore (A) in the mFARS strengthens the overall construct compared with the complete FARS.

Conclusions

A correlation-based psychometric analysis of the neurologic FARSn score justifies the overall validity of the scale. In addition, omission of items of limited functional significance as created in the mFARS improves the features of the measures. Such information is crucial to the ongoing application of the mFARS in natural history studies and clinical trials. Additional analyses of longitudinal changes will be necessary to fully ascertain its utility, especially in nonambulant patients.

From the Clinical Data Science GmbH (C.R.), Basel, Switzerland; Bruce Lefroy Centre for Genetic Health Research (L.A.C., M.B.D.), Murdoch Children's Research Institute, Parkville, Victoria, Australia; Department of Paediatrics (L.A.C., M.B.D.), University of Melbourne, Parkville, Victoria, Australia; Department of Neurology (S.H.S.), McKnight Brain Institute, Room, Gainesville, FL; University of Minnesota (K.B.); University of Chicago (C.M.G.); Ohio State University (J.C.H.); Divisions of Neurology and Clinical and Metabolic Genetics (G.Y.), Department of Paediatrics, the Hospital for Sick Children, University of Toronto, Ontario, Canada Hospital; University of Rochester (B.R.); University of Iowa (K.D.M.); Emory University (G.W.); University of South Florida (T.Z.); Friedreich's Ataxia Research Alliance (S.P.), Downingtown, PA; and Division of Neurology (D.R.L.), Children's Hospital of Philadelphia.

Go to Neurology.org/NG for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

ADL = activities of daily living; **CTT** = classical test theory; **FA-COMS** = Friedreich Ataxia–Clinical Outcome Measures Study; **FARS** = Friedreich Ataxia Rating Scale; **FARSn** = FARS neurologic examination; **FDS** = functional disability staging; **FRDA** = Friedreich ataxia; **ICARS** = International Cooperative Ataxia Rating Scale; **IRT** = Item Response Theory; **mFARS** = modified FARS; **PCA** = principal component analysis; **SARA** = Scale for the Assessment and Rating of Ataxia.

Inherited ataxias vary in clinical presentation, yet most do share cerebellar pathology leading to loss of balance and limb coordination, cardinal features that need to be measured to evaluate the efficacy of an intervention. Friedreich ataxia (FRDA) exemplifies this challenge of a complex neuropathology, including degeneration of the spinal cord, peripheral sensory nerves, and the dentate nucleus of the cerebellum,¹ which demands a broad but sensitive measure to capture the effect of neuroanatomical dysfunction on clinical progression of affected patients.

A first widely accepted such instrument was the International Cooperative Ataxia Rating Scale (ICARS),² but its use in clinical studies in FRDA^{3–7} has diminished^{8,9} in favor of advanced developments. One, the Scale for the Assessment and Rating of Ataxia (SARA)^{10,11} focuses on a compact structure and quick administration. Concurrently, the Friedreich Ataxia Rating Scale (FARS) combines timed measures of performance, functional disability staging (FDS), and a patient-reported outcome (activities of daily living [ADL]) with a paramount neurologic examination, the FARS neurologic examination (FARSn). Both SARA and FARS are now generally accepted and used in 2 large natural history studies in Europe^{12–15} and the United States.^{16,17}

The relationships of the FARSn with other rating scales,¹¹ patient-reported outcomes,^{18–20} disease duration and FDS,²¹ age at onset, and repeat length^{16,18,21–24} are well established. We summarize the revisions of the FARSn examination after its introduction¹⁸ and, based on classical test theory²⁵ (CTT), provide a psychometric analysis of the construct at its subscores, the lack of which has been acknowledged.^{11,26,27}

Methods

Standard protocol approvals, registrations, and patient consents

Informed consent was obtained from all participants before starting evaluations. The study was approved by the institutional review board at all participating sites.

Study design and participants

We used data from all participants recruited via the Collaborative Clinical Research Network in Friedreich Ataxia²⁸ into the Friedreich Ataxia–Clinical Outcome Measures Study (FA-COMS).¹⁷ Results from this ongoing and actively recruiting registry study have been published widely.^{16,22,24} Briefly, the FA-COMS has minimal inclusion/exclusion criteria (age 4–80 years

and clinical and genetic confirmation of FRDA), and the follow-up time now (as of 2019) is up to 15 years. Participant characteristics cover the full spectrum of FRDA with regard to age at disease onset, disease duration, and disease severity. There are 12 participating sites (number of participants): Children's Hospital of Philadelphia (417), University of California Los Angeles (191), Murdoch Children's Research Institute (179, Melbourne, Australia), Emory University (73), University of South Florida (36), University of Iowa (26), University of Florida (20), University of Chicago (20), Sick Kids Hospital (19, Toronto, Canada), University of Minnesota (17), Ohio State University (10), and University of Rochester (4).

Friedreich Ataxia Rating Scale

With the exception of the FARSn, the components of the FARS examination, ADL, timed measures, and FDS were not changed since its original publication.¹⁸ E.g., FARS disability staging (FDS) is derived from an ordinal score from 1 to 6 (no, minimal, mild, moderate, severe, and total disability), independent from FARS ratings. It is graded in units of 0.5 linked by descriptors to ambulation status and overall function (5 being permanent wheelchair user).¹⁸

The revision of the FARSn examination was initially driven by item reduction based on interrater reproducibility, and insertion of 2 items related to stance without visual aid.²¹ The resulting version (total of 125 points, figure 1) consists of 5 subscales directed to bulbar function (FARS A, maximum of 11 points), upper limb coordination (FARS B, 36), lower limb coordination (FARS C, 16), peripheral nervous system (FARS D, 26), and upright stability (FARS E, 36). This version is used in the FA-COMS¹⁷ since 2003 and has been used as an end point in clinical studies.^{29–31}

More recently, the need to focus solely on functional abilities led to a reduction in items, specifically in the peripheral nervous system and bulbar components.¹⁶ This resulted in the new, modified FARS (mFARS) score (mFARS, 93), which is used in currently ongoing and recently finished clinical trials.^{30–33} It is important that all items of the mFARS are included in the full FARSn examination; only the bulbar subscore (mA) was modified (now excluding items A1, facial atrophy and A2, and tongue atrophy), and the peripheral nervous system (D) subscore is now omitted. The remaining subscales, upper limb coordination (B), lower limb coordination (C), and upright stability (E), are identical in both the mFARS and the FARSn (see figure 1 for description and scoring for overall scales and individual subscales).

Table 1 Demographic characteristics of the cohort; percentages are based on the disease onset group

Mean (SD)	Disease onset group			Total
	<15 y (early)	15-24 y (intermediate)	>24 y (late)	
N (% ^a)	695 (68.7)	209 (20.7)	107 (10.6)	1,011
Male (%)	354 (50.9)	102 (48.8)	45 (42.1)	501 (49.6)
Age	19.41 (10.31)	32.32 (11.09)	50.12 (10.86)	25.33 (14.50)
Disease duration	10.95 (9.67)	14.40 (10.70)	15.18 (9.10)	12.11 (9.97)
Symptom onset	8.46 (3.36)	17.92 (2.54)	34.94 (9.38)	13.22 (9.41)
Age at diagnosis	12.15 (5.08)	22.96 (6.96)	41.82 (11.99)	17.47 (11.42)
GAA1 ^b	718.01 (209.29)	504.53 (197.76)	273.79 (194.32)	628.23 (252.91)
GAA2 ^c	941.51 (186.53)	852.80 (245.93)	795.34 (285.90)	908.24 (217.94)
Point mutation, N (%)	40 (5.8)	10 (4.8)	5 (4.7)	55 (5.4)
Disease stage				
1-2	319 (45.9)	80 (38.3)	32 (29.9)	431 (42.6)
3-4	145 (20.9)	77 (36.8)	51 (47.7)	273 (27.0)
5-6	231 (33.2)	52 (24.9)	24 (22.4)	307 (30.4)

^a Based on the overall cohort.

^b N = 939 (excluding point mutations).

^c N = 937.

(15–24 years), and 107 (10.6%) had late onset (≥ 25 years). In the 2 later-onset groups, fewer patients were in an early disease phase (stage 1–2, 38.3% and 29.9%), but there also were fewer patients in later, nonambulatory stages (stage 5–6, 24.9% and 22.4%, respectively). Overall, 1 third of participants (33.2% in the early-onset group, 30.4% of all participants) were already nonambulatory at their first visit. In line with previous reports, about 5.4% of the patients were carrying a point mutation. Because these patients show increased risk of atypical disease progression,^{40,41} they are typically excluded from clinical studies. However, the potential effect on the results was deemed minimal, and we included these patients in all analyses.

Psychometric properties of the FARS

Scaling characteristics of both full scores and all respective subscales are summarized in table 2, and the full corrected item-subscale correlations for FARSn subscores in table 3 (a full correlation table for mFARS is provided as table e-1, [links.lww.com/NXG/A189](https://www.links.lww.com/NXG/A189)). Mean corrected item-total correlations for subscales mA, B, C, D, and E, which are forming the mFARS, were 0.58 or above, whereas the A and the D subscale from FARSn showed mean values of 0.40 and 0.42, already indicating that the step from FARSn to the mFARS leads to more stable constructs. An additional benefit from the exclusion of A1 and A2 is the reduction of ceiling effects in the modified bulbar subscore (mA), leading to a less skewed distribution.

Cronbach α values were satisfactory for the upper limb (A, 0.87) and lower limb (B, 0.91), as well as for the upright

stability scale (E, 0.87). The modified bulbar subscale (mA) in the mFARS showed an α of 0.65, which is improved from the initial 4-item bulbar subscale (A) in FARSn, but still has to be considered a low value, clearly missing the suggested criterion of 0.9.^{42,43} For the initial FARSn subscores A and D, Cronbach α (0.53 and 0.56, respectively) shows that the modifications resulted in improved probabilities in mFARS. For the complete examinations, both the FARSn and the mFARS showed a Cronbach α of 0.92. Constructs covered by subscales of a total score should be related (intercorrelated), but also different to cover different aspects of a disease. A range of 0.30–0.70 has been suggested,⁹ and this condition was met for all subscores in both overall scales.

All subscales, as well as the complete scores, correlated with disease duration and FDS and, to a lesser extent, with age. Detailed item-own subscale and item-other subscale correlations (table 3) further corroborate construct validities of the subscores and full scales. All items in the mFARS subscales correlated with coefficients of >0.3 with their subscores, indicating that these items measure a common construct and collect a similar amount of information.^{9,34} In addition, only 1 item (E1, sitting posture) of the FARSn/mFARS examinations did not correlate higher with its own subscale than with other subscales. This item correlated to a similar extent with all 3 subscales (table 3) in mFARS, and also with the peripheral nervous system subscale D in FARSn. As a criterion for a margin between correlations between its own subscale and others, 2 SDs have been

Table 2 Scaling characteristics of sub- and total scores in the FARS neurologic examination, plus total FARSn and the mFARS^a

FARS subscale/items	mA—Bulbar (modified)	B—Upper limb	C—Lower limb	E—Upright stability	mFARS	FARSn	A—Bulbar ^b	D—Peripheral
No. of items	2	5	2	9	18	25	4	5
Corrected item–total correlations								
Mean	0.58	0.67	0.84	0.60	0.65	0.57	0.40	0.42
Range	0.55–0.61	0.41–0.77	0.84–0.84	0.24–0.86	0.25–0.86	0.17–0.86	0.22–0.60	0.17–0.72
Subscale characteristics								
Mean score (SD)	0.91 (0.90)	13.66 (6.97)	8.57 (4.73)	25.06 (8.32)	48.21 (18.52)	60.34 (22.17)	1.10 (1.08)	11.94 (4.64)
Observed score (max)	0–5	0–36	0–16	0–36	2–92	6–118	0–7	0–26
Floor (%)	33	2	2	0	0	0	25	0
Floor (10th percentile, %)	45	6	3	1	1	0	71	1
Ceiling (90th percentile, %)	0	1	22	31	2	1	0	2
Ceiling (%)	0	1	21	4	0	0	0	1
Skewness	1.02	0.52	0.40	–0.42	0.16	0.34	1.56	0.56
Cronbach α	0.65	0.87	0.91	0.87	0.92	0.92	0.53	0.56
Subscale intercorrelations								
Bulbar (modified)	—	0.63	0.57	0.55	0.68	0.67 ^b	—	0.49 ^b
Upper limb		—	0.68	0.66	0.88	0.86	0.64	0.57
Lower limb			—	0.76	0.88	0.88	0.56	0.70
Peripheral				—	—	0.81	0.49	—
Upright stability					0.92	0.91	0.54	0.93
Correlation with other parameters								
Disease duration	0.24	0.28	0.36	0.37	0.42	0.43	0.24	0.29
Age	0.10	0.07	0.09	0.09	0.10	0.10	0.09	0.04
Disability staging	0.35	0.41	0.56	0.76	0.73	0.72	0.33	0.38

Abbreviations: FARS = Friedreich Ataxia Rating Scale; mFARS = modified FARS.

^a Pearson correlation coefficients, corrected for overlap and scale reliability³⁷

^b Analyzed using FARSn subscales; a complete table of subscale correlations for mFARS subscales is provided in table e-1 (links.lww.com/NXG/A18).

suggested,^{9,34} which was fulfilled by all items but E1 (sitting posture) and E2B (stance, feet apart, with eyes closed), which also correlated well with subscale C (lower limb coordination). For the peripheral nervous system subscore in FARSn, except for D2 (muscle weakness), none of the items in the peripheral nervous system subscale correlated well with any other subscale (table 3). Remarkably, weaker item–subscale correlations were also found for items E3B (stance feet together, eyes closed), E4 (tandem stance), and E5 (stance on dominant foot) (see below).

Floor and ceiling effects

Ceiling effects were apparent in subscale C (heel-shin tap and heel-shin slide, possible range 0–8), with 21% of observations showing the maximum score. Notably, even after loss of ambulation, patients can perform the heel-shin tests (subscale C). For subscale E (upright stability), we found that item E1 (sitting position, possible range 0–4) is scored at 4 (indicating greatest impairment) in less than 5% of the observations, which masks total score/ceiling effects for subscale E. We also reported a ceiling effect at 90% of the maximum on this item

Table 3 Item-subscale correlations of the full FARSn^a

FARS subscale/items		A—Bulbar	B—Upper limb	C—Lower limb	D—Peripheral	E—Upright stability
Bulbar						
Facial atrophy ^b	A1	0.22	0.23	0.20	0.28	0.15
Tongue atrophy ^b	A2	0.35	0.25	0.18	0.23	0.17
Cough	A3	0.66	0.50	0.52	0.51	0.48
Speech	A4	0.70	0.64	0.54	0.47	0.51
Upper limb coordination						
Finger-finger	B1	0.45	0.52	0.31	0.34	0.29
Nose-finger	B2	0.69	0.82	0.63	0.60	0.57
Dysmetria	B3	0.76	0.85	0.67	0.62	0.64
Rapid movements	B4	0.60	0.79	0.59	0.55	0.62
Finger taps	B5	0.72	0.82	0.67	0.63	0.61
Lower limb coordination						
Heel-shin slide	C1	0.69	0.69	0.89	0.79	0.76
Heel-shin tap	C2	0.68	0.69	0.90	0.83	0.75
Peripheral nervous system^b						
Muscle atrophy	D1	0.35	0.30	0.43	0.53	0.32
Muscle weakness	D2	0.72	0.61	0.74	0.73	0.62
Vibratory sense	D3	0.22	0.20	0.33	0.37	0.34
Position sense	D4	0.29	0.38	0.45	0.54	0.46
Deep tendon reflexes	D5	0.04	0.16	0.12	0.20	0.16
Upright stability						
Sitting posture	E1	0.78	0.78	0.79	0.79	0.70
Stance, feet apart	E2A	0.64	0.60	0.78	0.76	0.83
Stance, feet apart, eyes closed	E2B	0.40	0.50	0.54	0.53	0.72
Stance, feet together	E3A	0.50	0.56	0.63	0.60	0.80
Stance, feet together, eyes closed	E3B	0.22	0.31	0.3	0.32	0.45
Tandem stance	E4	0.20	0.25	0.26	0.27	0.44
Stance on the dominant foot	E5	0.14	0.18	0.17	0.13	0.32
Tandem walk	E6	0.48	0.45	0.52	0.49	0.74
Gait	E7	0.69	0.65	0.81	0.77	0.90

Abbreviations: FARS = Friedreich Ataxia Rating Scale; mFARS = modified FARS.

^a Corrected for item overlap and scale reliability³⁷.

^b A1, A2, and the peripheral nervous system subscore are not included in the mFARS scores. A complete table of item-subscale correlations for mFARS is provided in table e-1 (links.lww.com/NXG/A18).

Item-own subscale correlations are emphasized in bold.

(table 2). Specifically, for the E (upright stability) subscale, the ceiling effect corresponded well to the number of non-ambulatory patients in our sample (32%, FDS > 4, table 1).

Floor effects were only visible in the bulbar subscale mA and (and bulbar/A in FARSn). The removal of A1 (facial atrophy) and A2

(tongue atrophy) reduced the ceiling effects to some extent, still 43% of patients had scored 10% or less of the total in the modified (mA) bulbar subscale. We also analyzed the ceiling effects on an item-based level. All individual item scores in mFARS (grouped by subscales) are presented in figure 2 as heat maps from lowest score (light blue) to maximum score (dark blue).

Figure 2 Heat map of individual items within the modified FARS

Data are grouped by subscales. Five blue hues (from light to dark) correspond to increasing item scores in steps of 20%. FARS = Friedreich Ataxia Rating Scale.

In general, the heat maps for items in subscales A, B, and C reflect nicely the broad and balanced stages of function loss in this population. Floor effects discussed above are visible in items A3 (cough) and A4 (speech) and relevant but still workable ceiling effects in the lower limb coordination subscale C. In the 9-item subscale E (upright stability), however, although favorable validity and consistency/reliability properties are achieved ($\alpha = 0.87$, table 2), only items E1 (sitting) and E7 (gait) and to some extent E6 (tandem walk) show balanced distributions. Specifically, all 6 stance-related items E2 to E5 reveal distinct bimodal distributions, which means that a patient can either perform the test well (score 0) or will be unable to complete the item rating a maximum score of 4. This is most evident in items E2A (stance, feet apart, eyes open), E2B (stance, feet apart, eyes closed), and E3A (stance, feet together, eyes open). For the remaining items, E3B (stance feet together, eyes closed), E4 (tandem stance), and E5 (stance dominant foot), it is also true, but only very few patients do not score the maximum.

A grouped analysis shows that only 15.8% of all patients ($n = 160$) do not score the maximum result in all E3B (stance, feet apart, eyes closed)/E4 (tandem stance)/E5 (stance on dominant foot) items, or, the other way around, 98.5% of patients ($n = 996$) had the maximum possible score in at least 1 of these 3 items. In the ambulatory sub-population ($FDS < 5$, $N = 704$, 69.6%), these proportions were 22.5% ($n = 159$) and 97.9% ($n = 689$).

Factor analysis

For principal component (PCA) and factor analysis, several solutions for FARS_n and mFARS were evaluated. Using the Kaiser criterion⁴⁴ (retaining factors with an eigenvalue >1), PCA found 5 factors for the complete FARS_n. A more detailed examination of the factor loading structure did not result in overall meaningful groupings. Especially, the items of the peripheral nervous system (D) subscore loaded into components together with items of other subscores (data not shown).

For mFARS, 3 principal components with eigenvalues >1 were found, and 1 additional with an eigenvalue of 0.83. Factor analysis using 4 factors resulted in a separation of independent, clinically meaningful factors. Items in FARS E (upright stability) and C (lower limb coordination) loaded strongest in component 1, upper limb coordination items (FARS B) loaded into component 2, and component 3 included solely the 2 bulbar function items of FARS A. The 3 remaining items of the FARS E (upright stability), with high amounts of “unable” (E3B, E4 and E5) loaded into the 4th component. Cumulatively, these 4 factors explained 70% of the variance in the overall construct.

Discussion

For our psychometric analysis of the FARS_n examination, we used the FA-COMS cohort, comprehensively representing the 2 most important dimensions of FRDA, being individuals from the complete severity spectrum of FRDA (as measured by age at symptom onset or GAA1/GAA2 repeat length), and at the same time including all disease durations. Moreover, the predominance of early vs late onset and children vs adults in recruitment is most likely representative of the incidence, but not necessarily prevalence of FRDA. Affected individuals progress fastest in the early phases of the disease²⁰; we used baseline data only to achieve optimal coverage of disease phases. This population profile should be ideal for the analyses of inherently sample-dependent psychometric properties of any scale. Still, overall correlation parameters will revolve around the proportion of early-/late-stage patients and the ratio of ambulant/nonambulant patients.

In the present study, classical psychometric correlation analysis (item-own vs item-other subscale correlations) confirms the validity and structure of the FARS_n examination, but specifically endorses the modifications leading to the mFARS. Overall, both scales showed appropriate item-subscale groupings, inter-subscale correlations, and internal consistency (Cronbach $\alpha > 0.90$). However, 2 sets of items within the complete 125-point FARS_n scale show clear weaknesses.

First, items in the peripheral nervous system (D) subscore itself have inefficient within subscale correlations and form a weak construct. This may reflect the arbitrary grouping of items within FARS D (peripheral nervous system). In particular, muscle strength and atrophy, items which have

a slower decline (and thus lower scoring) and may be secondary to both peripheral and CNS pathology, are grouped with deep tendon reflexes, which, being a hallmark feature of FRDA, are usually absent or diminished on first presentation therefore receiving a maximum score. Second, the A1 (facial atrophy)/A2 (tongue atrophy) items correlate weakly with other items in the bulbar (FARS A) subscore and other parts of the FARS examination. The remaining items in the bulbar subscore (related to cough and speech) have residual floor effects, yet they show considerable intercorrelation justifying their inclusion in the mFARS. Also, the Cronbach α value is improved for the modified bulbar (FARS mA) subscale.

These results justify the exclusion of these 2 groups of items (FARS D and A1/A2) in the newly formed modified mFARS score. Of interest, the requirement to focus on more patient-driven tasks initiated this change without knowledge of the psychometric properties of the items that quantify the functional ability of the individual with FRDA. As a result, intra-subscale correlations and reliability (Cronbach α) of the mFARS score are improved, whereas floor effects in FARS A1 (facial atrophy) and A2 (tongue atrophy) are avoided, demonstrating that the items reflecting functional ability match the most psychometrically robust in the mFARS. In addition, this shows that progression captured by the FARS examination(s) does not match the decline of dysfunctional sensory pathways in FRDA and that sensory loss is less relevant for the patient.

In addition, the present analysis demonstrates the detailed features of the FARS E (upright stability) subscale, which drives quantification of progression in the early phase of the disease. Its 6 stance-related items (E2A, B; E3A, B; E4, E5), are scored almost exclusively with extreme values (0 or the maximum, 4). This might indicate that in context of an otherwise slowly progressing condition, these abilities are lost rapidly. Alternatively, the extreme values could be a function of the item constructs. The easier, initial 3 stance items (E2A, E3A, and E2B) show preferable interitem correlations and should function well within the item response theory (IRT). On the other hand, the increasing difficulty of the remaining stance items results in very few patients in this cohort to perform E3B, E4, and E5. In most situations, these items thus contribute little to the total scores besides addition of noise. However, they also may provide useful information to the overall mFARS score in clinical studies that target those individuals with FRDA who are very early in the disease process. These results also indicate that all 6 stance items could be useful in classifying patients according to function and in longitudinal analysis of those individuals with the mildest of FRDA symptoms. The remaining 3 items in subscore E show beneficial psychometric properties; noteworthy, E7 (gait) correlates with all items in mFARS (and FARSn) and has the best overall correlation with the total mFARS score and in itself directly defines individual ambulatory ability. Another crucial item for functional ability, the sitting posture item (E1), showed the weakest inter-subscale correlation in FARS E (when disregarding E3B, E4, and E5), perhaps as scoring on this item can be improved by

scoliosis surgery or physiotherapy. Overall, although ceiling effects in FARS E (upright stability) match the nonambulatory fraction of the cohort, it contributes well to the overall FARSn/mFARS score in the ambulatory portion of the cohort.

Subscales FARS B (upper limb coordination) and C (lower limb coordination) showed beneficial psychometric properties for inclusion in the overall FARSn/mFARS. Of note, the finger-finger test item (B1) showed comparatively low item-subscale correlation, similar to that found in the ICARS analysis.⁹ Still, the remainder of these subscales provide useful contributions to the scale.

The use of IRT^{34,45} to evaluate rating scales in FRDA has been proposed.^{26,46} However, although Rasch-based models have been extended to polytomous⁴⁵ and continuous^{47,48} rating scales, they are not easily applied to scales with complex subscale/item structures like FARS and ICARS. Similarly, the assumption of unidimensionality in IRT-based models impedes their implementation.^{49,50}

Factor analysis for FARSn confirmed previous such findings for the FARSn examination, including the equivocal overlap between factor loadings and subscores.¹¹ For the mFARS, a 4-factor solution was the most clinically and statistically appropriate and PCA provided strong support of the subscore structure. This analysis additionally backs the exclusion of the FARS D and the 2 bulbar items and indicates that a lower limb component (FARS C plus E) and the upper limb portion (FARS B) within the mFARS have value as individual scales for separate evaluation.

In this context, the conceptual differences between a multi-subscore scale like the mFARS and a more compact scale (SARA) become obvious. The latter has been designed especially for rapid application and a strong central construct, leading to its unidimensionality.^{10,11} These features should improve interrater variability, highly important, e.g., in large multicenter studies. However, SARA might be limited in more complex studies. In contrast, the mFARS provides a more detailed evaluation of overall patient status and a more complex yet valid construct. In addition, subcomponents of the mFARS might separately support the evaluation of individual domains, e.g., upper and lower limb function, in specific circumstances. This might prove valuable when investigating temporal differences in loss of function in separate domains or when focusing on later stage, exclusively non-ambulatory patients. Although the complete FARS may require more time to complete, much of the extra information is collected in other instruments in typical clinical trials. Thus, overall, the different measures are relatively similar in practical administration, allowing investigators to select the best measure for the clinical approach being tested.

As future clinical trials in FRDA may focus on either subgroup, potentially targeted with specific therapeutic interventions, or more diverse cohorts with broader therapeutic

interventions, the present analysis demonstrates that the mFARS examination captures the complex features of FRDA, even in early stages of the disease.

Disclosure

Disclosures available: [Neurology.org/NG](https://www.neurology.org/NG).

Acknowledgment

The authors thank the coordinators of the Friedrich Ataxia Clinical Outcome Measures Study (FACOMS) for collection of patient visit data.

Study funding

This work was funded by the Friedrich's Ataxia Research alliance (FARA).

Publication history

Received by *Neurology: Genetics* June 7, 2019. Accepted in final form September 23, 2019.

Appendix Authors

Author	Location	Role	Contribution
Christian Rummey, PhD	Clinical Data Science GmbH, Basel	Author	Drafted the initial manuscript, revised the manuscript, performed data analysis, and conceived the study
Louise Corben, PhD	Murdoch Children's Research Institute	Author	Performed data collection and critical revision
Martin Delatycki, MD	Murdoch Children's Research Institute	Author	Performed data collection and critical revision
S.H. Subramony, MD	University of Florida	Author	Performed data collection and critical revision
Jennifer M. Farmer, MD	Friedreich's Ataxia Research Alliance	Author	Critical revision
David R. Lynch, MD, PhD	Children's Hospital of Philadelphia	Author	Performed data collection, supervised collaboration, and critical revision
Grace Yoon, MD	University of Toronto	Author	Performed data collection
Theresa Zesiewicz, MD	University of South Florida	Author	Performed data collection
Joseph Chad Hoyle, MD	Ohio State University	Author	Performed data collection
Katherine Mathews, MD	University of Iowa	Author	Performed data collection,

Appendix (continued)

Author	Location	Role	Contribution
George Wilmot, MD	Emory	Author	Performed data collection
Christopher M. Gomez, MD	University of Chicago	Author	Performed data collection
Khalaf Bushara, MD	University of Minnesota	Author	Performed data collection
Susan Perlman, MD	UCLA	Author	Performed data collection
Bernard Ravina, MD	University of Rochester	Author	Performed data collection and critical revision

References

- Koeppen AH, Mazurkiewicz JE. Friedrich ataxia: neuropathology revised. *J Neuropathol Exp Neurol* 2013;72:78–90.
- Trouillas P, Takayanagi T, Hallett M, et al. International cooperative ataxia rating scale for pharmacological assessment of the cerebellar syndrome. The ataxia neuropharmacology committee of the world federation of neurology. *J Neurol Sci* 1997; 145:205–211.
- Di Prospero NA, Baker A, Jeffries N, Fischbeck KH. Neurological effects of high-dose idebenone in patients with Friedrich's ataxia: a randomised, placebo-controlled trial. *Lancet Neurol* 2007;6:878–886.
- Lynch DR, Perlman SL, Meier T. A phase 3, double-blind, placebo-controlled trial of idebenone in Friedrich ataxia. *Arch Neurol* 2010;67:941–947.
- Meier T, Perlman SL, Rummey C, Coppard NJ, Lynch DR. Assessment of neurological efficacy of idebenone in pediatric patients with Friedrich's ataxia: data from a 6-month controlled study followed by a 12-month open-label extension study. *J Neurol* 2012;259:284–291.
- A study of efficacy, safety and tolerability of idebenone in the treatment of Friedrich's ataxia (FRDA) patients (MICONOS). Available at: clinicaltrials.gov/ct2/show/NCT00905268. Accessed January 4, 2019.
- Long-term safety and tolerability of idebenone in Friedrich's ataxia patients (MICONOS extension). Available at: clinicaltrials.gov/ct2/show/NCT00993967. Accessed January 4, 2019.
- Saute JAM, Donis KC, Serrano-Munuera C, et al. Ataxia rating scales—psychometric profiles, natural history and their application in clinical trials. *Cerebellum* 2012;11:488–504.
- Cano SJ, Hobart JC, Hart PE, Korlipara LVP, Schapira AHV, Cooper JM. International Cooperative Ataxia Rating Scale (ICARS): appropriate for studies of Friedrich's ataxia? *Mov Disord* 2005;20:1585–1591.
- Schmitz-Hübisch T, du Montcel ST, Baliko L, et al. Scale for the assessment and rating of ataxia: development of a new clinical scale. *Neurology* 2006;66:1717–1720.
- Bürk K, Mälzig U, Wolf S, et al. Comparison of three clinical rating scales in Friedrich ataxia (FRDA). *Mov Disord* 2009;24:1779–1784.
- Reetz K, Dogan I, Costa AS, et al. Biological and clinical characteristics of the European Friedrich's Ataxia Consortium for Translational Studies (EFACTS) cohort: a cross-sectional analysis of baseline data. *Lancet Neurol* 2015;14:174–182.
- Reetz K, Dogan I, Hilgers RD, et al. Progression characteristics of the European Friedrich's Ataxia Consortium for Translational Studies (EFACTS): a 2 year cohort study. *Lancet Neurol* 2016;15:1346–1354.
- Reetz K, Dogan I, Hohenfeld C, et al. Nonataxia symptoms in Friedrich ataxia: report from the registry of the European Friedrich's Ataxia Consortium for Translational Studies (EFACTS). *Neurology* 2018;91:e917–e930.
- Patient registry of the European Friedrich's Ataxia Consortium for Translational Studies (EFACTS). Available at: clinicaltrials.gov/ct2/show/NCT02069509. Accessed April 29, 2019.
- Patel M, Isaacs CJ, Seyer L, et al. Progression of Friedrich ataxia: quantitative characterization over 5 years. *Ann Clin Transl Neurol* 2016;3:684–694.
- FA Clinical Outcome Measures (FA-COMS). Available at: clinicaltrials.gov/ct2/show/NCT03090789. Accessed April 29, 2019.
- Subramony SH, May W, Lynch D, et al. Measuring Friedrich ataxia: interrater reliability of a neurologic rating scale. *Neurology* 2005;64:1261–1262.
- Fahey MC, Corben L, Collins V, Churchyard AJ, Delatycki MB. How is disease progress in Friedrich's ataxia best measured? A study of four rating scales. *J Neurol Neurosurg Psychiatry* 2006;78:411–413.
- Tai G, Corben LA, Gurrin L, et al. A study of up to 12 years of follow-up of Friedrich ataxia utilising four measurement tools. *J Neurol Neurosurg Psychiatry* 2015;86:660–666.

21. Lynch DR, Farmer JM, Tsou AY, et al. Measuring Friedreich ataxia: complementary features of examination and performance measures. *Neurology* 2006;66:1711–1716.
22. Friedman LS, Farmer JM, Perlman S, et al. Measuring the rate of progression in Friedreich ataxia: implications for clinical trial design. *Mov Disord* 2010;25:426–432.
23. Milne SC, Murphy A, Georgiou-Karistianis N, Yiu EM, Delatycki MB, Corben LA. Psychometric properties of outcome measures evaluating decline in gait in cerebellar ataxia: a systematic review. *Gait Posture* 2018;61:149–162.
24. Regner SR, Lagedrost SJ, Plappert T, et al. Analysis of echocardiograms in a large heterogeneous cohort of patients with Friedreich ataxia. *Am J Cardiol* 2012;109:401–405.
25. Spector P. *Summated Rating Scale Construction*. Newbury Park: SAGE Publications, Inc; 1992.
26. Delatycki MB. Evaluating the progression of Friedreich ataxia and its treatment. *J Neurol* 2009;256:36–41.
27. Marelli C, Figoni J, Charles P, et al. Annual change in Friedreich's ataxia evaluated by the Scale for the Assessment and Rating of Ataxia (SARA) is independent of disease severity. *Mov Disord* 2012;27:135–139.
28. FARA—Collaborative Clinical Research Network in Friedreich's Ataxia [online]. Available at: curefa.org/. Accessed April 29, 2019.
29. Marcotulli C, Fortuni S, Arcuri G, et al. GIFT-1, a phase IIa clinical trial to test the safety and efficacy of IFN γ administration in FRDA patients. *Neurol Sci* 2016;37:361–364.
30. Zesiewicz T, Heerinx F, De Jager R, et al. Randomized, clinical trial of RT001: early signals of efficacy in Friedreich's ataxia: signals of Efficacy in Friedreich's Ataxia. *Mov Disord* 2018;33:1000–1005.
31. Zesiewicz T, Salemi JL, Perlman S, et al. Double-blind, randomized and controlled trial of EPI-743 in Friedreich's ataxia. *Neurodegener Dis Manag* 2018;8:233–242.
32. Lynch DR, Farmer J, Hauser L, et al. Safety, pharmacodynamics, and potential benefit of omaveloxolone in Friedreich ataxia. *Ann Clin Transl Neurol* 2018;6:15–26.
33. Lynch DR, Hauser L, McCormick A, et al. Randomized, double-blind, placebo-controlled study of interferon- γ 1b in Friedreich ataxia. *Ann Clin Transl Neurol* 2019;6:546–553.
34. Hobart J, Cano S. Improving the evaluation of therapeutic interventions in multiple sclerosis: the role of new psychometric methods. *Health Technol Assess* 2009;13:1–177.
35. R Core Team. *R: A Language and Environment for Statistical Computing* [online]. Vienna: R Foundation for Statistical Computing; 2018. Available at: R-project.org/. Accessed April 29, 2019.
36. Wickham H. Tidyverse: easily install and load the “tidyverse” [online]. 2017. Available at: CRAN.R-project.org/package=tidyverse. Accessed April 29, 2019.
37. Revelle W. *Psych: Procedures for Psychological, Psychometric, and Personality Research* [online]. Evanston: Northwestern University; 2018. Available at: [//CRAN.R-project.org/package=psych](http://CRAN.R-project.org/package=psych).
38. Metz G, Coppard N, Cooper JM, et al. Rating disease progression of Friedreich's ataxia by the International Cooperative Ataxia Rating Scale: analysis of a 603-patient database. *Brain* 2013;136:259–268.
39. Rummey C. *Loss of Ambulation in the CCRN-FA Natural History Study*. Italy: Pisa; 2017.
40. Lynch DR, Deutsch EC, Wilson RB, Tennekoon G. Unanswered questions in Friedreich ataxia. *J Child Neurol* 2012;27:1223–1229.
41. Galea CA, Huq A, Lockhart PJ, et al. Compound heterozygous *FXN* mutations and clinical outcome in Friedreich ataxia: insights from Frataxin structure and function. *Ann Neurol* 2016;79:485–495.
42. Lance CE, Butts MM, Michels LC. The sources of four commonly reported cutoff criteria: what did they really say? *Organ Res Methods* 2006;9:202–220.
43. Nunnally JC. *Psychometric Theory* (2nd ed.). New York, NY: McGraw-Hill; 1978.
44. Kaiser HF. The application of Electronic computers to factor Analysis. *Educ Psychol Meas* 1960;20:141–151.
45. Rasch G. *On General Laws and the Meaning of Measurement in Psychology*. Copenhagen, Denmark: Danmarks pædagogiske Institut; 1961.
46. Lynch DR, Farmer JM, Wilson RL, Balcer LJ. Performance measures in Friedreich ataxia: potential utility as clinical outcome tools. *Mov Disord* 2005;20:777–782.
47. Andrich D. A rating formulation for ordered response categories. *Psychometrika* 1978;43:561–573.
48. Müller H. A Rasch model for continuous ratings. *Psychometrika* 1987;52:165–181.
49. Mair P, Hatzinger R. Extended Rasch modeling: The eRm package for the application of IRT models in R. *J Stat Softw* [online serial] 2007;20. Accessed at: jstatsoft.org/v20/i09/. Accessed August 2, 2019.
50. Hohensinn C. *pcIRT: an R package for polytomous and continuous Rasch models*. *J Stat Softw* [online serial] 2018;84. Accessed at: jstatsoft.org/v84/c02/. Accessed August 2, 2018.

Neurology[®] Genetics

Psychometric properties of the Friedreich Ataxia Rating Scale

Christian Rummey, Louise A. Corben, Martin B. Delatycki, et al.

Neurol Genet 2019;5;

DOI 10.1212/NXG.0000000000000371

This information is current as of October 29, 2019

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2376-7839.

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/5/6/371.full.html
References	This article cites 36 articles, 1 of which you can access for free at: http://ng.neurology.org/content/5/6/371.full.html##ref-list-1
Citations	This article has been cited by 5 HighWire-hosted articles: http://ng.neurology.org/content/5/6/371.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Clinical neurology examination http://ng.neurology.org/cgi/collection/clinical_neurology_examination Clinical trials Observational study (Cohort, Case control) http://ng.neurology.org/cgi/collection/clinical_trials_observational_study_cohort_case_control Gait disorders/ataxia http://ng.neurology.org/cgi/collection/gait_disorders_ataxia Mitochondrial disorders http://ng.neurology.org/cgi/collection/mitochondrial_disorders Spinocerebellar ataxia http://ng.neurology.org/cgi/collection/spinocerebellar_ataxia
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2376-7839.