Galloping tongue syndrome in a *PRRT2* mutation carrier

Dolores Vilas, MD, PhD, Anna Marcé-Grau, PhD, Alfons Macaya, PhD, Josep Valls-Solé, MD, PhD, and Eduard Tolosa, MD, PhD

Correspondence

Dr. Tolosa
etolosa@clinic.cat

Neurol Genet 2019;5:e377. doi:10.1212/NXG.0000000000000377

Abnormal lingual movements are not uncommon. A rare lingual syndrome consisting of involuntary wave-like lingual movements has been labeled as “galloping tongue.”¹ Herein, we report a patient with galloping tongue syndrome carrier of a mutation in the *PRRT2* gene.

MORE ONLINE

▶ **Video**

Clinical findings

A 17-year-old woman presented with involuntary tongue movements within the past years, appearing only with tongue protrusion. “Wavy” contractions of the tongue were evident during sustained tongue protrusion (video 1). The contractions developed immediately after sticking the tongue out, persisted during protrusion, and disappeared while at rest. The movements could not be disrupted, suppressed, or entrained with examiner-given rhythmic activities. Abrupt body movements, sleep deprivation, or coffee intake did not affect the abnormal movement. There was no history of exposure to neuroleptic medication or head trauma. The tongue contractions seemed to begin in the posterior midline region and progress in a wavy fashion to the anterior and lateral parts of the tongue. The results of routine hematologic and blood biochemistry including thyroid function tests, copper levels, and ceruloplasmin were normal. A urine toxicology screen did not reveal the presence of any drug of abuse. An electrophysiologic study with electrodes applied on the surface of the tongue showed no electrical activity while at rest. With tongue protrusion, the EMG activity was segmented in discharges of 200–300 ms duration and separated by brief periods of silence, with an approximate frequency of 3 per second (figure 1). Her brain MRI was normal. Because the movement disorder did not affect the patient’s orolingual function, no treatment was started.

The patient’s sister, a 14-year-old, presented with short episodes of choreo-dystonic movements of the left arm and neck that appeared when she started to run since the age of 10. These episodes usually last a few seconds and occur at least 10 times per day. Her consciousness level was normal during episodes. Her medical history included migraine. She was diagnosed as having paroxysmal kinesigenic dyskinesias (PKD). Treatment with carbamazepine led to a complete remission of the movements.

There was no family history of epilepsy or infantile convulsions. The patient’s mother had severe recurrent migraine headaches. Her maternal cousin had been diagnosed with PKD.

The patient, her 2 sisters (1 healthy and the other affected with PKD), and their mother were available for *PRRT2* mutation screen (figure 2). Direct sequencing identified a heterozygous truncating mutation in the proband, her mother, and the symptomatic sister: GRC37/hg19: chr16:29825025 GA/A, NM_145239: c.650delG, producing the amino acid change NP_660282.2: p.R217Qfs*12. This variant is not listed in public genomic databases, including ExAC (exac.broadinstitute.org/).

From the Parkinson’s Disease and Movement Disorders Unit (D.V., E.T.), Neurology Service, ICN, Hospital Clínic, University of Barcelona, Spain; Pediatric Neurology Research Group (A.M.-G., A.M.), Vall d’Hebron Institut de Recerca (VHIR), Universitat Autònoma de Barcelona, Spain; Electrophysiology Unit (J.V.-S.), Neurology Service, ICN, Hospital Clínic, University of Barcelona, Spain; Institut August Pi I Sunyer Biomedical Research Institute (IDIBAPS) (J.V.-S., E.T.).

Go to Neurology.org/NG for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Figure 1 Electromyographic activity of the tongue recorded with surface electrodes during tongue protrusion

Discharges of 200–300 ms duration and separated by brief periods of silence, with an approximate frequency of 3 per second were observed during tongue protrusion. Broken lines have been added to underline each EMG discharge.

Discussion

Herein, we describe a patient with “galloping tongue” syndrome who was positive for the p.R217Q fs*12 mutation in the *PRRT2* gene. Galloping tongue is an uncommon movement disorder.^{1–5} The characteristics of these lingual movements have been variably described as transverse contractions, twisting, or undulating movements.^{2–5} Although these movements were present with the tongue at rest in most of the reported patients, in some cases the movements were also observed during sustained action.

Our patient had a mutation in the *PRRT2* gene known to be associated with PKD.⁶ However, she did not have the typical paroxysmal dyskinesias seen in PKD. Her abnormal movements were triggered by action but were not paroxysmal because they were present as long as the voluntary movement motion persisted, resembling an action dystonia. The topography of dyskinesia, isolated to the tongue, was also uncharacteristic of PKD. Still, we believe that the *PRRT2* mutation could be linked to the patient’s movement disorder because it is action related and known causes of the disorder have been ruled out. Furthermore, the phenotype of *PRRT2* mutations is not limited to dystonia and chorea typical of PKD, but episodic ataxia and paroxysmal

Figure 2 *PRRT2* mutational analysis in a family with paroxysmal neurologic disorder

Right, *PRRT2* electropherograms: Top, wildtype sequence in the proband’s healthy sister; and bottom, guanine deletion at position 650 occurs at the end of a cytosine streak, a well-known hotspot in PKD. The variant results in a premature termination of transcription (p.R217Qfs*12). Left, family tree with various phenotypes indicated.

torticollis have also been described.⁷ Seizures and migraine seem to be also part of the clinical spectrum.⁶

The cause of the lingual movements in our patient remains uncertain. A functional movement disorder seemed unlikely because the movement disorder could not be disrupted, suppressed, or entrained with examiner maneuvers, which had no impact in the patient’s life, and there was no obvious gain for the patient. The study of similar cases and additional *PRRT2* families will eventually clarify whether the “galloping tongue” syndrome is indeed a previously unrecognized manifestation of *PRRT2* mutations.

Author contributions

D. Vilas and E. Tolosa: drafting/revising the manuscript and conceptualization of the study. A. Marcé-Grau and A. Macaya: genetic analysis and revising the manuscript. J. Vall-Solé: electrophysiologic study and revising the manuscript.

Study funding

No targeted funding reported.

Disclosure

Disclosures available: Neurology.org/NG.

Publication history

Received by *Neurology: Genetics* May 28, 2019. Accepted in final form October 9, 2019.

References

- Keane JR. Galloping tongue: post-traumatic, episodic, rhythmic movements. *Neurology* 1984;34:251–252.
- Saddichha BA, Talukdar D. Serpentine tongue syndrome associated with risperidone long-acting injection treatment. *J Clin Psychopharmacol* 2014;34:657–658.
- Nagappa M, Sinha S, Saini J, Bindu P, Taly A. Undulating tongue in Wilson’s disease. *Ann Indian Acad Neurol* 2014;17:225–226.

4. Prashantha DK. Serpentine tongue: a rare manifestation following initiation of levodopa therapy in a patient with Parkinson's disease. *Parkinsonism Relat Disord* 2009;15:718–719.
5. Sheehy SHC, Lawrence T, Thevathasan AW. Serpentine tongue. A lingual dyskinesia. *Neurology* 2008;70:e87.
6. Méneret A, Grabli D, Depienne C, et al. PRRT2 mutations: a major cause of paroxysmal kinesigenic dyskinesia in the European population. *Neurology* 2012;79:170–174.
7. Castelnovo G, Renard D, De Verdal M, Luc J, Thouvenot E, Riant F. Progressive ataxia related to PRRT2 gene mutation. *J Neurol Sci* 2016;367:220–221.

Neurology[®] Genetics

Galloping tongue syndrome in a *PRRT2* mutation carrier

Dolores Vilas, Anna Marcé-Grau, Alfons Macaya, et al.

Neurol Genet 2019;5;

DOI 10.1212/NXG.0000000000000377

This information is current as of November 12, 2019

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/5/6/e377.full.html
References	This article cites 7 articles, 0 of which you can access for free at: http://ng.neurology.org/content/5/6/e377.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://ng.neurology.org/cgi/collection/all_genetics All Movement Disorders http://ng.neurology.org/cgi/collection/all_movement_disorders
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.