

Aicardi-Goutières syndrome due to a paternal mosaic *IFIH1* mutation

Victoria Tüngler, MD, MSc, Marion Doeblner-Neumann, MD, Michaela Salandin, MD, Peter Kaufmann, MD, Christine Wolf, PhD, Nadja Lucas, PhD, Florian Harmuth, MSc, Jennifer Reichbauer, Ingeborg Krägeloh-Mann, MD, Rebecca Schüle, MD, and Min Ae Lee-Kirsch, MD

Correspondence

Dr. Lee-Kirsch
minae.lee-kirsch@
uniklinikum-dresden.de

Neurol Genet 2020;6:e384. doi:10.1212/NXG.0000000000000384

Progressive immune-mediated neurodegeneration is a central feature of Aicardi-Goutières syndrome (AGS), a monogenic disorder characterized by chronic activation of antiviral type I interferon (IFN).¹ Typically, AGS presents as subacute infancy-onset encephalopathy with microcephaly, leukodystrophy, and basal ganglia calcification, resulting in global developmental delay. AGS is either caused by loss-of-function mutations in *TREX1*, *RNASEH2B*, *RNASEH2C*, *RNASEH2A*, *SAMHD1*, or *ADAR*, encoding genes involved in the metabolism of nucleic acids, or by gain-of-function mutations in *IFIH1* encoding the cytosolic RNA sensor melanoma differentiation-associated protein 5 (MDAS).¹ The phenotypic spectrum of *IFIH1*-associated mutations includes intracerebral vasculopathy, bilateral striatal necrosis, and isolated spastic paraparesis.

We report the rare case of AGS due to paternal mosaicism for an *IFIH1* mutation in 2 brothers. The study was conducted with approval by the ethics committees of the University of Tübingen and Technische Universität Dresden, and written informed consent was obtained. Both siblings were born at term to healthy nonconsanguineous parents after uneventful pregnancies and with anthropometric birth data within normal limits. Their family history was unremarkable. After a period of normal development, both brothers presented with gait disturbances and progressive microcephaly. Bilateral lower limb spasticity manifested at the age of 18 months in the older brother (II:1) after he had learned to walk unsupported, whereas the younger brother (II:2) became symptomatic at the age of 12 months before learning to walk (figure, A). Apart from mild hypertonicity of the left arm and minor dysarthria in the older brother, neither of the 2 children showed signs of additional motor or cognitive deficits. Brain MRI revealed symmetric hyperintensities within the periventricular white matter in both brothers, with hypomyelination more pronounced in the older sibling (figure, B). Blood counts, inflammatory markers, and liver and renal function tests were unremarkable. Both siblings were clinically diagnosed with hereditary spastic paraplegia. Sequencing of 136 HSP-related genes (HaloPLEX hereditary spastic paraplegia Panel) identified a heterozygous variant of *IFIH1* (NM_022168: c.2336 G>A, p. R779H) in both children. Of interest, the variant was also observed at low abundance in the blood-derived DNA sample of the clinically asymptomatic father. Sanger sequencing confirmed the heterozygous R779H variant in both children, while a weak mutation peak was also observed in the sequence pherogram of the father, confirming that he was mosaic for R779H (figure, C). Thus, both children inherited the R779H mutation through a germline mosaic from the father.


R779H has previously been reported in at least 8 patients with AGS occurring either as a dominant mutation with reduced penetrance or as de novo mutation.² We therefore investigated the family for signs of constitutive type I IFN activation in blood. Consistent with AGS, both brothers exhibited a strong IFN signature (IFN score $1,031.19 \pm 350.19$ in I:1 and 648.21 ± 219.61 in I:2, mean \pm SEM; normal range < 12.49). Although the mother showed no signs of IFN activation (IFN score 1.29), the father was also found to have an IFN signature (IFN score 404.04),

From the Department of Pediatrics (V.T., C.W., N.L., M.A.L.-K.), Medizinische Fakultät Carl Gustav Carus, Technische Universität Dresden; Department of Neuropediatrics (M.D.-N., I.K.-M.), University of Tübingen, Germany; Child Neurology and Psychiatry Unit (M.S.), Paediatric Department, Bolzano Regional Hospital; Child Haematology and Oncology Unit (P.K.), Paediatric Department, Bolzano Regional Hospital, Italy; Institute of Medical Genetics and Applied Genomics (F.H.), University of Tübingen; and Center for Neurology and Hertie-Institute for Clinical Brain Research (J.R., R.S.), University of Tübingen and German Center of Neurodegenerative Diseases, Germany.

Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by DFG.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.



(A) Pedigree. (B) MRI of II:1 at 2 years, delayed myelination and white matter hyperintensity (left, white arrows), and unchanged at 5 years (right). (C) Heterozygous *IFIH1* mutations in the children, weak mutation peak in the father (reverse sequence). (D) IFN scores ($p \leq 0.0007$, before vs after ruxolitinib), calculated as described.³ IFN = interferon.

consistent with the mosaic state of the R779H variant in his blood. Further examination of the father did not reveal microcephaly, vasculitis, or lupus-like symptoms. His blood counts and renal and liver function tests were unremarkable.

Uncontrolled activation of the MDA5 receptor because of activating *IFIH1* mutations results in constitutive type I IFN signaling.² Given the disease progression and lack of approved therapeutic options, we initiated off-label treatment with the Janus kinase (JAK) 1/2 inhibitor ruxolitinib, which inhibits downstream signaling at the IFN- α/β receptor. Ruxolitinib started at 5 and 7 years, respectively, with a dose of 0.5 mg/kg was well tolerated without any hematologic or infectious adverse events. Ruxolitinib was increased to 0.75 mg/kg over time. Both children responded with a significant reduction of the IFN signature (figure, D). The parents reported a marked improvement in their children's quality of life during ruxolitinib treatment, who were described to be less fatigued and to engage more motivated in physical activities. Improved concentration of the older brother had a positive effect on academic achievements. Both children were able to maintain and even moderately improve their motor abilities, with a progress more noticeable in the younger brother, whose gait using orthoses improved by 40% after 8 months of treatment, as revealed by the dimension "walking, running, and jumping" of the Gross Motor Function Measure.

Clinical improvement observed in the patients supports previous reports, indicating that JAK inhibition may be therapeutically effective in type I IFN-driven disorders.^{3–7} Timely diagnosis is of clinical importance because early therapeutic intervention may modify the course of the disease and prevent further neurologic damage. Our findings also suggest that parental germline mosaicism may be more common than previously presumed in patients with AGS with apparent *de novo* *IFIH1* mutation with significant implications for genetic counseling.

Acknowledgment

The authors are thankful to the family for participation in this study. The authors thank Diana Federl and Kerstin Engel for excellent technical assistance.

Study funding

Supported by grants from the Deutsche Forschungsgemeinschaft (LE1074/4-1 and grant 369799452/404459235 to ML-K and TU421/1-2 to VT), the NEUROMICS network (F5–2012–305121 to RS), Horizon 2020 'Solve-RD' (grant 779257 to RS), the National Institute of Health (NIH) (grant 5R01NS072248 to RS) and the Bundesministerium für Bildung und Forschung via funding for the TreatHSP consortium (01GM1905 to RS and MD-N).

Disclosure

Disclosures available: [Neurology.org/NG](https://www.neurology.org/NG).

Publication history

Received by *Neurology: Genetics* June 7, 2019. Accepted in final form October 31, 2019.

Appendix Authors

Name	Location	Role	Contribution
Victoria Tüngler, MD, MSc	Technische Universität Dresden, Germany	Author	Study design, clinical assessment, analysis, and interpretation of the data, and drafting of the manuscript
Marion Doeblner-Neumann, MD	University of Tübingen, Germany	Author	Clinical assessment and acquisition and analysis of the data
Michaela Salandin, MD	Regional Hospital of Bolzano, Italy	Author	Clinical assessment and acquisition of the data
Peter Kaufmann, MD	Regional Hospital of Bolzano, Italy	Author	Clinical assessment and acquisition of the data
Christine Wolf, PhD	Technische Universität Dresden, Germany	Author	Acquisition and analysis of the data
Nadja Lucas, PhD	Technische Universität Dresden, Germany	Author	Acquisition and analysis of the data

Appendix (continued)

Name	Location	Role	Contribution
Florian Harmuth, MSc	University of Tübingen, Germany	Author	Acquisition and analysis of the data
Jennifer Reichbauer	University of Tübingen, Germany	Author	Acquisition and analysis of the data
Ingeborg Krägeloh-Mann, MD	University of Tübingen, Germany	Author	Clinical assessment, analysis, and interpretation of the data
Rebecca Schüle, MD	University of Tübingen, Germany	Author	Study design and analysis and interpretation of the data
Min Ae Lee-Kirsch, MD	Technische Universität Dresden, Germany	Author	Study design, clinical assessment, analysis, and interpretation of the data, and drafting of the manuscript

References

1. Lee-Kirsch MA. The type I interferonopathies. *Annu Rev Med* 2017;68:297–315.
2. Rice GI, Del Toro Duany Y, Jenkinson EM, et al. Gain-of-function mutations in IFIH1 cause a spectrum of human disease phenotypes associated with upregulated type I interferon signaling. *Nat Genet* 2014;46:503–509.
3. König N, Fiehn C, Wolf C, et al. Familial chilblain lupus due to a gain-of-function mutation in STING. *Ann Rheum Dis* 2017;76:468–472.
4. Tüngler V, König N, Günther C, et al. Response to: “JAK inhibition in STING-associated interferonopathy” by Crow et al. *Ann Rheum Dis* 2016;75:e76.
5. Kim H, Brooks KM, Tang CC, et al. Pharmacokinetics, pharmacodynamics, and proposed dosing of the oral JAK1 and JAK2 inhibitor baricitinib in pediatric and young adult CANDLE and SAVI patients. *Clin Pharmacol Ther* 2018;104:364–373.
6. Kothur K, Bandodkar S, Chu S, et al. An open-label trial of JAK 1/2 blockade in progressive IFIH1-associated neuroinflammation. *Neurology* 2018;90:289–291.
7. Zimmermann N, Wolf C, Schwenke R, et al. Assessment of clinical response to Janus kinase inhibition in patients with familial chilblain lupus and TREX1 mutation. *JAMA Dermatol* 2019;155:342–346.

Neurology[®] Genetics

Aicardi-Goutières syndrome due to a paternal mosaic *IFIH1* mutation

Victoria Tüngler, Marion Doebler-Neumann, Michaela Salandin, et al.

Neurol Genet 2020;6;

DOI 10.1212/NXG.0000000000000384

This information is current as of December 19, 2019

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/6/1/e384.full.html
References	This article cites 7 articles, 2 of which you can access for free at: http://ng.neurology.org/content/6/1/e384.full.html#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://ng.neurology.org/cgi/collection/all_clinical_neurology All Genetics http://ng.neurology.org/cgi/collection/all_genetics All Movement Disorders http://ng.neurology.org/cgi/collection/all_movement_disorders All Pediatric http://ng.neurology.org/cgi/collection/all_pediatric
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.

