Association of a structural variant within the *SQSTM1* gene with amyotrophic lateral sclerosis

Julia Pytte, BSc (Hons), Ryan S. Anderton, PhD, Loren L. Flynn, PhD, Frances Theunissen, MBIomedSc, Leanne Jiang, BSc (Hons), Ianthe Pitout, PhD, Ian James, PhD, Frank L. Mastaglia, MD, Ann M. Saunders, PhD, Richard Bedlack, MD, PhD, Teepu Siddique, MD, Nailah Siddique, RN, Msn, and P. Anthony Akkari, PhD

Correspondence

Prof. Akkari
Anthony.akkari@perron.uwa.edu.au

Neurol Genet 2020;6:e406. doi:10.1212/NXG.0000000000000406

Abstract

Objective

As structural variations may underpin susceptibility to complex neurodegenerative diseases such as amyotrophic lateral sclerosis (ALS), the objective of this study was to investigate a structural variant (SV) within sequestosome 1 (*SQSTM1*).

Methods

A candidate insertion/deletion variant within intron 5 of the *SQSTM1* gene was identified using a previously established SV evaluation algorithm and chosen according to its subsequent theoretical effect on gene expression. The variant was systematically assessed through PCR, polyacrylamide gel fractionation, Sanger sequencing, and reverse transcriptase PCR.

Results

A reliable and robust assay confirmed the polymorphic nature of this variant and that the variant may influence *SQSTM1* transcript levels. In a North American cohort of patients with familial ALS (fALS) and sporadic ALS (sALS) ($n = 403$) and age-matched healthy controls ($n = 562$), we subsequently showed that the *SQSTM1* variant is associated with fALS ($p = 0.0036$), particularly in familial superoxide dismutase 1 mutation positive patients ($p = 0.0005$), but not with patients with sALS ($p = 0.97$).

Conclusions

This disease association highlights the importance and implications of further investigation into SVs that may provide new targets for cohort stratification and therapeutic development.

From the University of Western Australia (J.P., R.S.A., L.L.F., F.T., L.J., F.L.M., P.A.A.), Centre for Neuromuscular and Neurological Disorders, Crawley; Perron Institute for Neurological and Translational Science (J.P., R.S.A., L.L.F., F.T., L.J., I.P., F.L.M., P.A.A.), Nedlands; University of Notre Dame Australia (R.S.A.), School of Health Sciences; University of Notre Dame Australia (R.S.A.), Institute for Health Research, Fremantle; Murdoch University (L.L.F., I.P., P.A.A.), Centre for Molecular Medicine and Innovative Therapeutics; Murdoch University, Institute for Immunology and Infectious Diseases (I.J.), Western Australia, Australia; Department of Neurology (R.B.), Duke University School of Medicine, Durham, NC; Zinfandel Pharmaceuticals (A.M.S.), Inc.; Duke University (R.B.), ALS Clinic, Durham, NC; and Departments of Neurology, Pathology and Cell and Molecular Biology (T.S., N.S.), Northwestern University Feinberg School of Medicine, the Les Turner ALS Center and the Northwestern University Interdepartmental Neuroscience Program, Chicago, IL.

Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

ALS = amyotrophic lateral sclerosis; **CI** = confidence interval; **fALS** = familial ALS; **GWAS** = genome-wide association study; **I/D** = insertion/deletion; **mRNA** = messenger RNA; **ONS** = olfactory neurosphere derived; **OR** = odds ratio; **sALS** = sporadic ALS; **SOD1** = superoxide dismutase 1; **SQSTM1** = sequestosome 1; **RT-PCR** = reverse transcriptase PCR; **SV** = structural variant.

A growing body of literature indicates that structural variants (SVs) are important in complex diseases because their role in gene expression and messenger RNA (mRNA) regulation is now emerging.^{1–4} Given the potential implication of SVs in neurodegenerative disorders,^{5–7} an investigation of SVs within associated genes may provide insight toward the pathogenic mechanisms involved and enable the identification of new therapeutic targets.

Mutations in sequestosome 1 (*SQSTM1*), and aggregation of the *SQSTM1*-encoded p62 protein, have been identified in patients with amyotrophic lateral sclerosis (ALS) and other neurodegenerative diseases.^{8–13} P62 is a multifunctional protein that binds ubiquitin and is involved in autophagy, proteasomal degradation of ubiquitinated proteins, mitophagy, and cellular signaling.^{14–16} Variants within *SQSTM1* or surrounding *SQSTM1* may contribute to the diverse presentation observed between the patients with ALS. How mutations in *SQSTM1* influence ALS is not fully understood; however, it is believed that structural changes to the protein may affect adapter function of *SQSTM1* to the LC3 protein in the nascent autophagosome and impair autophagy of proteins that are not recyclable by the proteasome.¹⁷ Other cellular systems may also be affected such as ubiquitin binding and regulation of cellular processes including DNA repair, endocytosis, and signal transduction.^{8,14}

In the current study, a potential influential insertion/deletion (I/D) within intron 5 of *SQSTM1* was identified using an *in silico* short SV evaluation algorithm.¹⁸ Reverse transcriptase PCR (RT-PCR) analysis revealed a link between the I/D and altered transcript levels. An exploratory association study in a sporadic and familial North American ALS cohort established an association with familial ALS (fALS), particularly superoxide dismutase 1 (*SOD1*) mutation-positive patients but not sporadic ALS (sALS). We further examined whether this SV was associated with age of disease onset and duration in *SOD1* mutation-positive patients and found no statistically significant association.

Methods

Standard protocol approvals, registrations, and patient consents

This study was approved by the ethical standards of the relevant institutional review board, the Human Research Ethics Committee of the University of Western Australia (RA/4/20/5308). Participants were enrolled after informed consent was obtained. Clinical data were collected according to the Health

Insurance Portability and Accountability Act Standards of Confidentiality and Disclosure and approved by the Northwestern University institutional review board (STU0012722/CR5_STU00012722) and the Duke University institutional review board (PRO00040665/323682). All patients were diagnosed by board-certified neurologists and met the revised El Escorial World Federation of Neurology criteria for diagnosis with ALS.¹⁹

SV identification

An SV evaluation algorithm was used to locate the variant within the gene, *SQSTM1*.¹⁸ Detailed methods are described in the supplementary material (see e-methods, links.lww.com/NXG/A229).

PCR amplification and Sanger sequencing

Detailed methods are described in the supplementary material. Sequences of all primers used for this study are outlined in table e-1 (links.lww.com/NXG/A229).

Olfactory neurosphere-derived cells

Culturing information can be found in the supplementary data.

RT-PCR and densitometry

Detailed methods are described in the supplementary material. Sequences of all primers used for this study are detailed in table e-1 (links.lww.com/NXG/A229).

ALS and healthy control participants

A cohort of 196 fALS, 207 sALS, and 562 healthy control participants were recruited into the Neurologic Diseases Registry, Northwestern University, Chicago, USA, and Duke University, North Carolina, USA. The 196 fALS cases from Northwestern University belong to 74 different families with possible within-family genetic correlations. The mutation data for all fALS patients are detailed in table e-2 (links.lww.com/NXG/A229).

Statistics

Data are reported as mean $\pm$ SD where appropriate. Statistical differences in genotype proportions for independent cases were assessed using Pearson χ^2 test with Yates correction. Proportions and ages of onset involving familial groups were analyzed using the mixed effects regression models to account for possible within-family correlations. Durations were analyzed using Cox proportional hazards models with clustering to account for the correlations. Analyses were carried out in IBM SPSS Statistics version 25.0 (IBM Co., Armonk, NY) and R version 3.4.3 (R Foundation for Statistical Computing, Vienna, Austria).

Data availability

Study data for the primary analyses presented in this report are available on reasonable request from the corresponding author.

Results

Identification of a polymorphic variant within the *SQSTM1* gene

An SV evaluation algorithm was used to identify the genetic variant located within *SQSTM1*. The identified cytosine adenine adenine adenine (CAAA) I/D within intron 5 of *SQSTM1* was evaluated on the NCBI database; although the region had been previously reported (NC_0000005.0), no disease associations were established. The recorded NCBI data demonstrated uncertainty to its polymorphic nature; currently, the region is recorded as a series of insertion/deletions of varying length, with no allelic frequency data. Conventional polyacrylamide gel electrophoresis revealed a biallelic variant, consisting of a heterozygous I/D genotype in lanes 2, 3, and 7, a homozygous D/D genotype in lanes 4 and 8, and a homozygous I/I genotype in lanes 5, 6, and 9 (figure 1A). Sanger sequencing confirmed that the I allele contained a CAAA insertion when compared with the D allele (figure 1B). To determine the effect of the CAAA SV on *SQSTM1* expression, *SQSTM1* was assessed in a panel (n = 3) of olfactory neurosphere-derived (ONS) cells obtained from the healthy controls. Semiquantitative RT-PCR revealed a stepwise increase in the level of *SQSTM1* transcript, with the I/I genotype yielding the lowest levels and the I/D and D/D genotypes producing a 1.77-fold and 2.47-fold relative increase in *SQSTM1* transcript, respectively (figure 1C).

The *SQSTM1* variant is associated with fALS

Following the identification of the CAAA I/D variant, we proceeded to determine the variant frequency in a cohort of 196 fALS patients, 207 sALS patients, and 562 healthy age-matched controls. The characteristics of the study participants are summarized in table 1, including sex, age, disease duration, and family history of each cohort. Of the ALS cohort, patients with familial mutations (48.6%) were further grouped by affected gene, specifically *C9orf72* (2.5% of patients with ALS), *SOD1* (41.4% of patients with ALS), or *TDP-43* (4.7% of patients with ALS).

There was no difference in the frequency of the *SQSTM1* variant observed between healthy controls and sALS cases ($\chi^2 = 0.032$, df = 2, $p = 0.984$, odds ratio [OR] = 0.974, 95% confidence interval [CI]: 0.660, 1.436). However, there was a difference in *SQSTM1* variant frequency observed between healthy controls and fALS patients ($\chi^2 = 12.791$, df = 2, $p = 0.002$, OR = 1.79, 95% CI: 1.250, 2.562 assuming independence of cases), with the homozygous (I/I) genotype over-represented in the fALS cohort (33.2%) when compared with the controls (21.7%). When analyzed using mixed effects regression models to account for possible within-family correlations, the (I/I) association remained both genotypical and allelic ($p = 0.013$ and $p = 0.0036$, respectively, table 2). This

Figure 1 Characterization of the *SQSTM1* variant

(A) PCR and native polyacrylamide gel electrophoresis across a random selection of control DNA samples and compared against a 100bp ladder. (B) Sanger sequencing of the I allele and D allele. (C) *SQSTM1* transcript levels of *SQSTM1* exon 4-7 analyzed using RT-PCR on RNA from a panel of control ONS cells. Relative densitometry was calculated with *SQSTM1* transcript signal standardized to each respective GAPDH signal. GAPDH = glyceraldehyde 3-phosphate dehydrogenase; I/D = insertion/deletion; ONS = olfactory neurosphere derived cells; RT-PCR = Reverse transcriptase PCR; *SQSTM1* = sequestosome 1.

SQSTM1 association was stronger when examining the subset of fALS patients carrying mutations in *SOD1* ($\chi^2 = 16.754$, df = 2, $p = 0.0002$, OR = 1.869, 95% CI: 1.281, 2.726 assuming independent cases). When accounting for family structures, the *SQSTM1* variant was strongly associated with patients carrying a *SOD1* mutation both genotypically and allelically ($p = 0.001$ and $p = 0.0005$, respectively) (table 2). When analyzed with *SOD1*-ASV mutation-positive patients excluded, the strong *SOD1* association remains ($\chi^2 = 6.801$, $p < 0.009$, OR = 1.830, 95% CI: 1.157, 2.896).

The *SQSTM1* variant is not associated with age at onset of disease or survival in patients with *SOD1*.

To determine if the *SQSTM1* variant is associated with *SOD1* mutation-positive ALS patient outcomes, age at onset and survival were analyzed. There was no association observed between the age at onset in patients with the I/I (Mean Rank = 87.25), I/D (Mean Rank = 76.40), or D/D (Mean Rank =

Table 1 Characteristics of patients and healthy control participants

Variable	Mean (SD) or n	
	Patients (n = 403)	Controls (n = 562)
Sex		
Male	208	296
Female	195	266
Age (y)	52.38 (12.99)	49.37 (12.56)
Disease duration (mo)	43.21 (46.98)	—
Family history		—
Sporadic	207	
Familial	196	—

84.43) genotypes ($p = 0.2$, mixed model regression, figure 2A). Survival curves were generated to compare durations of each genotypic group (figure 2B). No association was observed between patients carrying each genotype and their survival after taking into account of familial correlations via a clustered Cox regression ($p = 0.55$).

Discussion

A growing body of literature has highlighted SVs, their abundance throughout the human genome, and their potential role in the pathogenesis of ALS and other neurodegenerative diseases.^{20–24} SVs are responsible for greater diversity at the nucleotide level between 2 human genomes than any other form of genetic variations and are three-fold more likely to associate with genome-wide association studies (GWASs) signals than single nucleotide polymorphisms (SNPs).²⁰ SVs that remain cryptic to current sequencing algorithms are likely to account for disease-causing variation in unsolved Mendelian disorders and missing heritability in complex disorders.^{20,23} SVs may affect gene expression and therefore may play an important but understudied role in disease susceptibility.^{20,25} Recent discoveries of SVs as informative disease risk markers for rare genetic disorders provide compelling evidence for ongoing investigation into the association between SVs and rare genetic diseases.^{26–30} The novel bioinformatics SV evaluation algorithm tool prioritizes potential functional/causal SVs within candidate regions identified using GWAS.¹⁸ In ALS and all rare genetic diseases, these highly polymorphic markers are often overlooked, largely because of the limitations in the current gene sequencing platforms such as next-generation sequencing and GWAS, which are primarily designed to detect SNPs.

Two direct lines of evidence link *SQSTM1* to ALS: the first through etiology on account of mutations associated with ALS and second through pathology, where p62 immune-reactive

Table 2 Association of *SQSTM1* variant and ALS disease status

sALS	Control n (%)	sALS n (%)	p Value
<i>SQSTM1</i> CAAA variant			
I/I	122 (21.7)	44 (21.3)	0.98
I/D	292 (52.0)	109 (52.7)	
D/D	148 (26.3)	54 (26.1)	
I allele	536	197	0.97
D allele	588	217	
Familial ALS	Control n (%)	fALS n (%)	p Value ^a
<i>SQSTM1</i> CAAA variant			
I/I	122 (21.7)	65 (33.2)	0.0013
I/D	292 (52.0)	97 (49.5)	0.50
D/D	148 (26.3)	34 (17.3)	0.08
I allele	536	227	0.0036
D allele	588	165	
<i>SOD1</i> mutation positive	Control n (%)	<i>SOD1</i> n (%)	p Value ^a
<i>SQSTM1</i> CAAA variant			
I/I	122 (21.7)	57 (34.1)	0.001
I/D	292 (52.0)	87 (52.1)	0.99
D/D	147 (26.3)	23 (13.8)	0.014
I allele	536	201	0.0005
D allele	588	133	

Abbreviations: ALS = amyotrophic lateral sclerosis; fALS = familial ALS; I/D = insertion/deletion; sALS = sporadic ALS; *SQSTM1* = sequestosome 1.
^a Each row compared using random effects to account for familial correlation.

inclusions are found in virtually all forms of ALS and ALS-frontotemporal dementia.⁸ We now provide a third link of *SQSTM1* variants to the fALS. *SQSTM1*-encoded p62 protein is a key scaffold involved in cellular signaling and protein degradation through the autophagosome-lysosome system.¹⁷ *SQSTM1*/p62 mutations may confer a toxic gain of function through protein interactions, leading to dysregulation of cell signaling pathways, protein misfolding, and aggregation.⁸ Evidence for this in the literature suggests parallels between p62 and other proteins associated with neurodegeneration, including *SOD1*-linked ALS, *C9orf72*, ubiquitin 2, TDP-43, FUS, optineurin, beta-amyloid, α -synuclein, and tau.^{8,31} Owing to its role in protein degradation, overexpression of p62 has been shown to be protective in some neurodegenerative animal models, but overexpression in a *SOD1* ALS model was found to accelerate disease onset.^{32–34} Taken together, these studies suggest that a fine balance in p62 levels is required for optimal signaling and protein clearance. Consequently, small changes in

Figure 2 Association of the *SQSTM1* variant with age at onset of disease and survival

(A) The median and distribution of age at onset (years) of *SOD1* mutation-positive patients with ALS ($n = 167$) grouped by each *SQSTM1* variant genotype. (B) Kaplan-Meier survival curves of *SOD1* mutation-positive patients, comparing the *SQSTM1* genotypes assuming independent measurements. A robust log-rank test accounting for familial correlation was performed to assess any association between the groups. Survival was measured in months from ALS diagnosis until death. ALS = amyotrophic lateral sclerosis; *SOD1* = superoxide dismutase 1; *SQSTM1* = sequestosome 1.

the level of p62 expression, which may be produced by variants such as the *SQSTM1* intron 5 I/D, could tip the balance of p62 expression, contributing to the disease. Previously identified mutations within *SQSTM1* have been associated with ALS.^{35,36} As such, it has been suggested that these mutations may have a direct role in ALS pathogenesis, presenting as an important target for future therapy.⁸

In this study, we hypothesized that structural variations in *SQSTM1* may uncover novel susceptibility factors that underpin this disease. *SQSTM1* mutations, and p62

aggregation, have been reported not only in ALS but also in other complex neurodegenerative diseases, including Parkinson disease, Paget disease of bone, Alzheimer disease and frontotemporal dementia.^{8–13} A semiquantitative assessment of the mRNA expression revealed a difference in *SQSTM1* transcript levels in the ONS cells carrying each genotype. Neural tissue derived from olfactory mucosa, such as primary ONS cells, provide informative cellular models for neurodegenerative diseases. Primary ONS cells are neural progenitor cells and more accurately reflect motor neuron cellular function than fibroblasts and PBMCs, and unlike postmortem brain tissue, they can be isolated from living patients.^{37,38} As such, the altered *SQSTM1* transcript levels observed in the ONS cells with the I/I genotype may be reflective of altered *SQSTM1* expression in motor neurons. Although this result requires validation in a larger sample size, it does suggest some level of regulation by the SV on *SQSTM1* transcript expression. This may be due to altered transcription efficiency, splicing, or transcript stability, potentially translating to altered protein expression that may contribute to disease pathologies.

An evaluation of the *SQSTM1* variant within a cohort of 403 patients with ALS and 562 healthy age-matched controls revealed an association with fALS disease risk, but not age of disease onset or disease severity. Replications in additional fALS cohorts are necessary to truly elucidate the nature of this variant. Causality has been reported between ALS and *SQSTM1* variants rs796051870, rs776749939, rs796052214, and as such, the ongoing investigation between ALS and *SQSTM1* gene variants is necessary.^{8,39,40} P62 has previously been reported as overexpressed and accumulated in inclusions of sporadic inclusion body myositis, reinforcing its importance in neurodegenerative diseases.¹² We anticipate that as additional SVs are identified, these will further stratify other relevant disease phenotypes such as age at onset, duration, and disease progression.

Acknowledgment

The authors are deeply indebted to the late Allen D. Roses, PhD, who provided mentorship to several of the authors and who originally conceptualized and initiated this research. The authors would also like to thank the contribution of Prof. Alan Mackay Sim and Prof. George Mellick (Griffith University, Institute for Drug Discovery) for the generous gift of the ONS cell cultures. The study was in part funded by NIH grants to TS (NS050641, NS046535), the Les Turner ALS Foundation/Herbert C. Wenske Foundation Professorship and the Les Turner ALS Research and Patient Care Center.

Study funding

The study was funded by research support from the Perron Institute for Neurologic and Translational Science and the Les Turner Foundation for ALS. The funders have no role in the design of the study and collection, analysis, decision to publish, interpretation of data, or preparation of the manuscript.

Disclosure

Disclosures available: [Neurology.org/NG](https://www.neurology.org/NG).

Publication history

Received by *Neurology: Genetics* July 19, 2019. Accepted in final form January 23, 2020.

Appendix Authors

Name	Location	Contribution
Julia Pytte, BSc (Hons)	The Perron Institute for Neurological and Translational Science, Nedlands, Australia	Acquisition of the data, analysis and interpretation, statistical analysis, and critical revision of the manuscript for important intellectual content
Ryan S. Anderton, PhD	The Perron Institute for Neurological and Translational Science, Nedlands, Australia	Study concept and design, statistical analysis, study supervision, and critical revision of the manuscript for important intellectual content
Loren L. Flynn, PhD	Murdoch University, Murdoch, Australia	Study concept and design, analysis and interpretation, study supervision, and critical revision of the manuscript for important intellectual content
Frances Theunissen, MBIomedSc	The Perron Institute for Neurological and Translational Science, Nedlands, Australia	Analysis and interpretation and critical revision of the manuscript for important intellectual content
Leanne Jiang, BSc (Hons)	The Perron Institute for Neurological and Translational Science, Nedlands, Australia	Analysis and interpretation and critical revision of the manuscript for important intellectual content
Ianthe Pitout, PhD	The Perron Institute for Neurological and Translational Science, Nedlands, Australia	Study concept and design and acquisition of the data
Ian James, PhD	Murdoch University, Murdoch, Australia	Statistical analysis and critical revision of the manuscript for important intellectual content
Frank Mastaglia, MD	The Perron Institute for Neurological and Translational Science, Nedlands, Australia	Critical revision of the manuscript for important intellectual content
Ann M. Saunders, PhD	Duke University, Durham, NC	Study concept and design and critical revision of the manuscript for important intellectual content

Appendix (continued)

Name	Location	Contribution
Richard Bedlack, MD, PhD	Duke University, Durham, NC	Study concept and design, acquisition of the data, and critical revision of the manuscript for important intellectual content
Teepu Siddique, MD, DSc (hc), FAAN	Northwestern University, Evanston, IL	Study concept and design, acquisition of the data, and critical revision of the manuscript for important intellectual content
Nailah Siddique, RN, Msn	Northwestern University, Evanston, IL	Study concept and design and critical revision of the manuscript for important intellectual content
P. Anthony Akkari, PhD	The Perron Institute for Neurological and Translational Science, Crawley, Australia	Study concept and design, acquisition of data, analysis and interpretation, and critical revision of the manuscript for important intellectual content and study supervision

References

1. Chakravarti A, Kapoor A. Mendelian puzzles. *Science* 2012;335:930–931.
2. Feuk L, Carson AR, Scherer SW. Structural variation in the human genome. *Nat Rev Genet* 2006;7:85–97.
3. Stankiewicz P, Lupski JR. Structural variation in the human genome and its role in disease. *Annu Rev Med* 2010;61:437–455.
4. Lupski JR. Structural variation mutagenesis of the human genome: impact on disease and evolution. *Environ Mol Mutagen* 2015;56:419–436.
5. Lyall DM, Royle NA, Harris SE, et al. Alzheimer's disease susceptibility genes APOE and TOMM40, and hippocampal volumes in the Lothian birth cohort 1936. *PLoS One* 2013;8:e80513.
6. Lutz MW, Saul R, Linnertz C, et al. A cytosine-thymine (CT)-rich haplotype in intron 4 of SNCA confers risk for Lewy body pathology in Alzheimer's disease and affects SNCA expression. *Alzheimers Dement* 2015;11:1133–1143.
7. Linnertz C, Anderson L, Gottschalk W, et al. The cis-regulatory effect of an Alzheimer's disease-associated poly-T locus on expression of TOMM40 and apolipoprotein E genes. *Alzheimers Dement* 2014;10:541–551.
8. Fecto F, Yan J, Vemula SP, et al. SQSTM1 mutations in familial and sporadic amyotrophic lateral sclerosis. *Arch Neurol* 2011;68:1440–1446.
9. Rea SL, Majcher V, Searle MS, Layfield R. SQSTM1 mutations—bridging paget disease of bone and ALS/FTLD. *Exp Cell Res* 2014;325:27–37.
10. Zatloukal K, Stumpfner C, Fuchsichler A, et al. p62 Is a common component of cytoplasmic inclusions in protein aggregation diseases. *Am J Pathol* 2002;160:255–263.
11. Lowe J, Blanchard A, Morrell K, et al. Ubiquitin is a common factor in intermediate filament inclusion bodies of diverse type in man, including those of Parkinson's disease, Pick's disease, and Alzheimer's disease, as well as Rosenthal fibres in cerebellar astrocytomas, cytoplasmic bodies in muscle, and mallory bodies in alcoholic liver disease. *J Pathol* 1988;155:9–15.
12. Nogalska A, Chiara AE, Ae T, et al. p62/SQSTM1 is overexpressed and prominently accumulated in inclusions of sporadic inclusion-body myositis muscle fibers, and can help differentiating it from polymyositis and dermatomyositis. *Acta Neuropathol* 2009;118:407–413.
13. Kwok CT, Morris A, de Belleruche JS. Sequestosome-1 (SQSTM1) sequence variants in ALS cases in the UK: prevalence and coexistence of SQSTM1 mutations in ALS kindred with PDB. *Eur J Hum Genet* 2014;22:492–496.
14. Kirkin V, McEwan DG, Novak I, Dikic I. A Role for ubiquitin in selective autophagy. *Mol Cell* 2009;34:259–269.
15. Welchman RL, Gordon C, Mayer RJ. Ubiquitin and ubiquitin-like proteins as multifunctional signals. *Nat Rev Mol Cell Biol* 2005;6:599–609.

16. Rogov V, Dötsch V, Johansen T, Kirkin V. Interactions between autophagy receptors and ubiquitin-like proteins form the molecular basis for selective autophagy. *Mol Cell* 2014;53:167–178.
17. Pankiv S, Clausen TH, Lamark T, et al. p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem* 2007;282:24131–24145.
18. Saul R, Lutz MW, Burns DK, Roses AD, Chiba-Falek O. The SSV evaluation system: a tool to prioritize short structural variants for studies of possible regulatory and causal variants. *Hum Mutat* 2016;37:877–883.
19. Brooks BR, Miller RG, Swash M, Munsat TL, World Federation of Neurology Research Group on Motor Neuron Diseases. El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Mot Neuron Disord* 2000;1:293–299.
20. Chaisson MJP, Sanders AD, Zhao X, et al. Multi-platform discovery of haplotype-resolved structural variation in human genomes. *Nat Commun* 2019;10:1784.
21. Sudmant PH, Rausch T, Gardner EJ, et al. An integrated map of structural variation in 2,504 human genomes. *Nature* 2015;526:75–81.
22. Roses AD. Short sequence variations (SSVs) may explain multiplicity of SOD1-SNP mutations in Amyotrophic Lateral Sclerosis. Presented at the 66th Annual Meeting American Society of Human Genetics; October 19, 2016; Vancouver.
23. Roses AD, Akkari PA, Chiba-Falek O, et al. Structural variants can be more informative for disease diagnostics, prognostics and translation than current SNP mapping and exon sequencing. *Expert Opin Drug Metab Toxicol* 2016;12:135–147.
24. Frazer KA, Murray SS, Schork NJ, Topol EJ. Human genetic variation and its contribution to complex traits. *Nat Rev Genet* 2009;10:241–251.
25. Li YC, Korol AB, Fahima T, Nevo E. Microsatellites within genes: structure, function, and evolution. *Mol Biol Evol* 2004;21:991–1007.
26. De Jesus-Hernandez M, Mackenzie IR, Boeve BF, et al. Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-Linked FTD and ALS. *Neuron* 2011;72:245–256.
27. Gijselinck I, Van Langenhove T, van der Zee J, et al. A C9orf72 promoter repeat expansion in a Flanders-Belgian cohort with disorders of the frontotemporal lobar degeneration-amyotrophic lateral sclerosis spectrum: a gene identification study. *Lancet Neurol* 2012;11:54–65.
28. Renton AE, Majounie E, Waite A, et al. A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 2011;72:257–268.
29. Sproviero W, Shatunov A, Stahl D, et al. ATXN2 trinucleotide repeat length correlates with risk of ALS. *Neurobiol Aging* 2017;51:178.e-1–178.
30. Roses AD, Lutz MW, Amrine-Madsen H, et al. A TOMM40 variable-length polymorphism predicts the age of late-onset Alzheimer's disease. *Pharmacogenomics J* 2010;10:375–384.
31. Deng HX, Chen W, Hong ST, et al. Mutations in UBQLN2 cause dominant X-linked juvenile and adult-onset ALS and ALS/dementia. *Nature* 2011;477:211–215.
32. Doi H, Adachi H, Katsuno M, et al. p62/SQSTM1 differentially removes the toxic mutant androgen receptor via autophagy and inclusion formation in a spinal and bulbar muscular atrophy mouse model. *J Neurosci* 2013;33:7710–7727.
33. Hadano S, Mitsui S, Pan L, et al. Functional links between SQSTM1 and ALS2 in the pathogenesis of ALS: cumulative impact on the protection against mutant SOD1-mediated motor dysfunction in mice. *Hum Mol Genet* 2016;25:3321–3340.
34. Mitsui S, Otomo A, Nozaki M, et al. Systemic overexpression of SQSTM1/p62 accelerates disease onset in a SOD1 H46R-expressing ALS mouse model. *Mol Brain* 2018;11:1–16.
35. Gal J, Strom A-L, Kwinter DM, et al. Sequestosome 1/p62 links familial ALS mutant SOD1 to LC3 via a ubiquitin-independent mechanism. *J Neurochem* 2009;111:1062–1073.
36. Zhou L, Wang H, Ren H, et al. Bcl-2-dependent upregulation of autophagy by sequestosome 1/p62 in vitro. *Acta Pharmacol Sin* 2013;34:651–656.
37. Féron F, Perry C, Girard SD, Mackay-Sim A. Isolation of adult stem cells from the human olfactory mucosa. In: Reynolds B, Deleyrolle L, editors. *Methods in Molecular Biology*. New Jersey: Humana Press; 2013:107–113.
38. Matigian N, Abrahamsen G, Sutharsan R, et al. Disease-specific, neurosphere-derived cells as models for brain disorders. *Dis Model Mech* 2010;3:785–798.
39. Teyssou E, Takeda T, Lebon V, et al. Mutations in SQSTM1 encoding p62 in amyotrophic lateral sclerosis: genetics and neuropathology. *Acta Neuropathol* 2013;125:511–522.
40. Le Ber I, Camuzat A, Guerreiro R, et al. SQSTM1 mutations in French patients with frontotemporal dementia or frontotemporal dementia with amyotrophic lateral sclerosis. *JAMA Neurol* 2013;70:1403–1410.

Neurology[®] Genetics

Association of a structural variant within the *SQSTM1* gene with amyotrophic lateral sclerosis

Julia Pytte, Ryan S. Anderton, Loren L. Flynn, et al.

Neurol Genet 2020;6;

DOI 10.1212/NXG.0000000000000406

This information is current as of February 27, 2020

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2376-7839.

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/6/2/e406.full.html
References	This article cites 38 articles, 4 of which you can access for free at: http://ng.neurology.org/content/6/2/e406.full.html##ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://ng.neurology.org/content/6/2/e406.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://ng.neurology.org/cgi/collection/all_genetics All Neuromuscular Disease http://ng.neurology.org/cgi/collection/all_neuromuscular_disease Amyotrophic lateral sclerosis http://ng.neurology.org/cgi/collection/amyotrophic_lateral_sclerosis_ Association studies in genetics http://ng.neurology.org/cgi/collection/association_studies_in_genetics
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2376-7839.