Molecular Diagnosis in 100% of Dystrophinopathies

Are We There Yet?

Elena Pegoraro, MD, PhD

Neurol Genet 2021;7:e529. doi:10.1212/NXG.0000000000000529

Correspondence

Dr. Pegoraro
elena.pegoraro@unipd.it

Duchenne muscular dystrophy (DMD) is experiencing its renaissance. After being the first muscular dystrophy in which the disease gene has been identified, it is now the first muscular dystrophy to be treated, in some cases, with personalized, dystrophin-restoring therapy.¹ In spite of these tremendous advancements in the treatment of DMD, in this issue of *Neurology® Genetics*, Waddel et al.² represent the challenges in achieving a genetic diagnosis in a cohort of male patients with elevated serum creatine kinase, dystrophin protein studies suggesting an underlying dystrophinopathy, yet normal multiplex ligation-dependent probe amplification and exomic sequencing.

The authors combined omic and bioinformatic analyses to identify causative mutations in their small cohort of undiagnosed patients. This is an important, well-designed, and timely study, which through intelligent use of updated omic resources and experimental data addresses several important issues in the field of dystrophinopathies.

First, the importance to achieve a molecular diagnosis in any suspected dystrophinopathy to allow patients to access potential therapies and families to receive appropriate genetic counselling. Although most of the currently available therapies to restore dystrophin expression (bioactive molecules that modulate the translation machinery reading through premature stop codons during messenger RNA [mRNA] translation, or treating RNA expression with anti-sense oligonucleotides) are not amenable to this cohort of patients harboring large structural rearrangements of the *DMD* gene, the foreseen availability of gene therapy potentially targeted to any DMD patients, regardless of the *DMD* mutation is emerging and likely it will deeply impact prognosis.^{1,3} Under these premises, the achievement of the molecular diagnosis in DMD is of a foremost relevance. The experimental approach used by Waddel et al.² demonstrates also the relevance of muscle biopsy in selected cases of unsolved muscular dystrophy. *DMD* mRNA isolated from skeletal muscle tissue will allow the localization of sequence anomalies in the transcript pointing to deep intronic causative mutations. Indeed, only the complementary use of genome sequencing, transcriptomics, and dystrophin protein studies have allowed a successful mutation identification in this cohort of patients.

Second, the authors attempted to define the amount of full-length dystrophin capable to rescue clinical phenotype using the remnant levels of normally spliced dystrophin *DMD* mRNA in patients with complex splicing mutations. The *vexata quaestio* “how much dystrophin is enough?” is now receiving a new answer.

It is well known that dystrophin is the strongest genetic modifier of the phenotype in the clinical spectrum of dystrophinopathies. If everybody agrees that “the more the better,” more controversies emerge to define the minimum amount necessary to be clinically relevant. Preclinical and single patient reports suggest that 15%–20% homogeneous dystrophin expression is sufficient to completely protect against eccentric contraction-induced injury, but still the debate is active.^{4,5} Waddel et al. using skeletal muscle biopsies reveals that normal dystrophin level $\sim 15 \pm 2\%$ correlates to the

RELATED ARTICLE

WGS and RNA Studies
Diagnose Noncoding
DMD Variants in Males
With High Creatine Kinase
Page e554

From the ERN Neuromuscular Unit, Department of Neurosciences, DNS, University of Padova, Padova, Italy.

Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures. Funding information is provided at the end of the article.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

milder spectrum of dystrophinopathies (i.e., myalgia without overt weakness), $\sim 10 \pm 2\%$ levels of dystrophin predicts a Becker muscular dystrophy phenotype with mild weakness and cardiac involvement, and 0%–5% levels of dystrophin results in severe Becker muscular dystrophy or DMD.² These predictions are in line with early studies in which level of dystrophin 29%–57% of normal was shown to be sufficient to avoid muscle weakness in X-linked dilated cardiomyopathy families.⁶ The novelty in the approach by Waddell et al is the estimate of normal dystrophin vs the amount of internally deleted dystrophin able to modulate the phenotype.¹ In the latter case, not only the amount but also the deleted region of the dystrophin gene is relevant to predict phenotype. Indeed, different regions in dystrophin have variable significance for its function, and thus, the phenotype prediction remain more complex and likely less precise. Upcoming clinical studies in the field of gene therapy will show how this “qualitative” factor will interact with sheer dystrophin quantity to determine the efficacy of various “microdystrophins” engineered into AAV vectors.

This study has impact for its translational implications in the emerging field of personalized medicine in dystrophinopathies

and will help assessing the efficacy of dystrophin-restoring therapies.

Study Funding

The author reports no targeted funding.

Disclosure

No relevant disclosures. Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosure.

References

1. Verhaart IEC, Aartsma-Rus A. Therapeutic developments for Duchenne muscular dystrophy. *Nat Rev Neurol* 2019;15:373–386.
2. Waddell LB, Breyer SJ, Cummings BB, et al. WGS and RNA studies diagnose non-coding DMD variants in males with high creatine kinase. *Neurol Genet* 2021;7:e554. doi:10.1212/NXG.0000000000000554.
3. Mendell JR, Sahenk Z, Lehman K, et al. Assessment of systemic delivery of rAAVrh74.MHCK7.micro-dystrophin in children with Duchenne muscular dystrophy. A non randomized controlled trial. *JAMA Neurol* 2020;77:1–10.
4. Godfrey C, Muses S, McClorey G, et al. How much dystrophin is enough: the physiological consequences of different levels of dystrophin in the mdx mouse. *Hum Mol Genet* 2015;24:4225–4237.
5. Todeschini A, Gualandi F, Trabaneli C, et al. Becker muscular dystrophy due to an intronic splicing mutation inducing a dual dystrophin transcript. *Neuromuscul Disord* 2016;26:662–665.
6. Neri M, Torelli S, Brown S, et al. Dystrophin levels as low as 30% are sufficient to avoid muscular dystrophy in the human. *Neuromuscul Disord* 2007;17:913–918.

Neurology[®] Genetics

Molecular Diagnosis in 100% of Dystrophinopathies: Are We There Yet?

Elena Pegoraro

Neurol Genet 2021;7;

DOI 10.1212/NXG.0000000000000529

This information is current as of January 29, 2021

Updated Information & Services	including high resolution figures, can be found at: http://ng.neurology.org/content/7/1/e529.full.html
References	This article cites 6 articles, 1 of which you can access for free at: http://ng.neurology.org/content/7/1/e529.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://ng.neurology.org/cgi/collection/all_genetics All Neuromuscular Disease http://ng.neurology.org/cgi/collection/all_neuromuscular_disease
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://ng.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://ng.neurology.org/misc/addir.xhtml#reprintsus

Neurol Genet is an official journal of the American Academy of Neurology. Published since April 2015, it is an open-access, online-only, continuous publication journal. Copyright © 2021 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2376-7839.